

A review of taxonomic studies of Arbuscular Mycorrhizal Fungi in Iran

E. Sedaghati[✉]

Department of Plant Protection, Faculty of Agricultural Sciences, Vali-e-Asr University of Rafsanjan, Rafsanjan Iran

M. Yazdanpanah

Department of Plant Protection, College of Plant Production, Gorgan University of Agricultural Sciences and Natural Resources, Gorgan, Iran

M. Nadi

Pistachio Research Center, Horticulture Sciences Research Institute, Agriculture Research Education and Extension Organization (AREEO), Rafsanjan, Iran

Abstract: Arbuscular Mycorrhizal Fungi (AMF) are one of the most essential beneficial soil microorganisms that can form a mutualistic symbiotic relationship with plants. AMF receive carbon from the host plant to complete their life cycle. In return, these fungi have copious roles for the host plants, including plant protection against pathogens, increasing abiotic stresses tolerance (drought and salinity), and enhancing water and mineral nutrient acquisition. Formerly, AMF was identified and classified merely based on morphological features of fungal spores such as mode of formation, wall structure, and subtending hyphal characteristics. Later on, molecular procedures were incorporated for AMF identification. PCR-based techniques led to the direct identification of AMF species that existed in plant roots or the rhizosphere. Several primers have been developed to increase the accuracy of AMF identification. Nowadays, classification systems of AMF are based on both morphological and molecular techniques. This paper aims to review the researches on the identification of AMF in Iran. Identification of AMF has received increasing interest over the past few decades. Authors have used a combination of morphological characteristics and DNA-based techniques to the identification of AMF. So far, more than 115 AMF species belonging to 22 genera have been identified and reported from different regions and plant communities such as croplands, forests, grasslands, etc. Understanding the community composition and diversity of AMF is vital for using them as biofertilizer in agriculture to reduce chemical inputs and increase sustainable crop production.

KEYWORDS: Community composition, Diversity, *Glomus*, Spore morphology, Symbiosis.

INTRODUCTION

Mycorrhiza is a symbiotic association between the plant root systems and soilborne fungi (Willis et al., 2013). Among the different types of mycorrhizae, arbuscular mycorrhizal fungi (AMF) are the most common symbiotic relationship with plants. AMF are obligate biotrophs of over 80 % of vascular plant species, which play a crucial role in nutrient cycling processes and maintaining ecosystem stability (Smith & Smith 2011). These fungi have an enormous ecological and economic impacts. The extra-radical hyphae of AMF, which are much thinner than fine roots, penetrate unreachable small spaces of soil and enhance the root surface area (Zou et al., 2021). Subsequently, they improve the growth of host plants and alleviate the effects of biotic and abiotic stresses (Chen et al., 2018, Zhang et al., 2020). AMF symbiosis enhances the growth and nutrition of host plants and reduces the need to fertilizers (Balestrini & Lumini 2018).

AMF species belonging to various genera have been reported from diverse habitats of the world, e.g. meadows, frigid regions, tropical regions, croplands, grasslands, tropical to temperate forests, alpine, dunes, deserts, etc. (Lovelock et al., 2003, Al-Yahya'ei et al., 2011, Varga et al., 2015). It should be noted that a small number of AMF species are limited to specific environmental conditions (Rosendahl 2008). The composition, abundance, and diversity of AMF are affected by many factors, like seasonal variations, edaphic characteristics, plant species, etc. (Bencherif et al., 2016, Turrini et al., 2017, Wang et al., 2021).

Morphological characteristics of spores are a basis for the identification of AMF species. The identification and classification of AMF based on morphological differences of the spores is complicated and frequently misleading or impossible and require a lot of information and experience (Kruger et al., 2012). Most of the recent molecular works have focused on using PCR-based techniques to analyze target sequences within the ribosomal DNA (rDNA) genes. This region is very suitable for designing general and specific primers due to having coded/conserved genes (SSU, 5S, and LSU) and non-coded / variable genes (ITS and IGS) (Symanczik 2016). A part of the large subunit (LSU)

or small subunit (SSU) rDNA gene or combination of both with internal transcribed spacer (ITS) have been chosen as a target for the Polymerase Chain Reaction (PCR) in many studies. These regions contain enough variation to distinguish and differentiate species of AMF (Binet et al., 2011, Johnson et al., 2016, Nielsen et al., 2016). Each of these regions alone is not enough for species resolution of AMF.

DNA is extracted from a certain amount of host plant roots or one spore or multispore or soil sample and is amplified using the AMF-specific primers (Ferrol & Lanfranco 2020). Until now, many researchers designed specific primers that would amplify all known AMF species. Molecular identification methods have the potential to study AMF ecology because they allow to identification and evaluation of the AMF taxa in root samples without the need for fungal spores. They also have the potential to identify AMF hyphae in the soil (Sanders 2004). So, the choice of the DNA region used for amplification is critically essential (Gorzalak et al., 2012).

The inability to accurately and adequately identify AMF species is caused by the nature of their obligate symbiont and the failure to independently culture any of them (Berruti et al., 2016). Identification and classification of the AMF are made based on morphological characteristics of spores and molecular tools applied separately or combined (Kruger et al., 2012, Wetzel et al., 2014). By far, more than 340 AMF species have been identified and described worldwide, and the

taxonomy of this fungal group is constantly being developed. The molecular techniques and morphological observations are shown that number of AMF species is greater than the present estimations and is expected up to 1600 (Tedersoo et al., 2018).

Isolation and identification of AMF species in each region are crucial steps prior to producing of fungal inoculum and employment in plantations and afforestation sites. In this regard, it is essential to know which types of AMF have symbiotic relationships with native plant species (Smith & Smith 2011). This review provides an overview of the morphological and molecular identification of AMF species in Iran over the past few decades.

Arbuscular Mycorrhiza Fungi

Mycorrhizal fungi are the most widespread and abundant fungi in nature and form a symbiosis relationship with many plants. The fungi belong to Ascomycota, Glomeromycota, Basidiomycota, and Mucoromycota (Smith & Read 2008). Mycorrhizal associations based on the type of fungus, host plant lineages, and structures produced by the root-fungus combination are classified into seven groups: endomycorrhiza (AM), ectomycorrhiza (EC), ectendomycorrhiza, ericoid, arbutoid, orchid and monotropoid (Tao et al., 2016) (Fig. 1). The movement of carbon from plant to fungus is common in all types of mycorrhizal association (Allen et al., 2003). Most often, mycorrhizal fungi are beneficial for host plants by improving the nutritional status of the plant (Chen et al., 2018).




Fig. 1. Schematic representation of various types of mycorrhizal fungi (Amballa & Bhumi 2017).

The arbuscular mycorrhizal symbiosis is probably the most widely beneficial interaction between fungi belonging to the Glomeromycota phylum and the host plant (Giovannini et al., 2020). AMF are obligate symbionts that able to establish a mutualistic symbiotic association with the majority (upwards of 80%) of vascular plants, including Bryophytes, Hepatics, Pteridophytes, Gymnosperms and Angiosperms, in terrestrial ecosystems (Gao et al., 2019, Zhang et al., 2020). AMF receives the carbohydrates and lipids necessary for the survival of the photosynthetically produced carbohydrates by the host plant; Such that 10 to 40 % of the carbohydrates can be absorbed by fungi (Begum et al., 2019, Genre et al., 2020). In return, AMF improve the acquisition of essential mineral nutrients (particularly phosphorus and nitrogen) and water supply. This improvement is done using development a branching mycelial network in the soil that can be expanded the host root surface area by 40 times (Hayashi et al., 2018, Zhang et al., 2019). Fundamentally, these fungi provide other benefits such as carbon recycling, improvement soil aggregation, production of plant growth hormones, enhance the photosynthetic efficiency of the host, and helping the plant to tolerance abiotic (such as drought, salinity, and heavy metals contamination)

and biotic stresses (pathogens and insects) through the implementation of various mechanisms (Ferrol et al. 2019, Lakmali et al., 2019, Li et al., 2020, Wu et al., 2021). As obligate biotrophs, AMF cannot complete their life stages without a compatible and living host plant (Smith & Read 2008). Under optimal conditions, an AMF spore (s) germinates, and after receiving signals released by the symbiotic partner, the hyphal germ tube navigates toward the host plant root (Tahat & Sijam 2012). Once in contact with the surface of the host's root, fungal hyphae form attachment structures, called "hyphopodia or appressoria" (Bonfante & Genre 2015). Then, hyphae penetrate toward the inner cortex cell and produce intraradical tree-like branched hyphae structures called "arbuscules". Arbuscules are the inorganic minerals (especially phosphorus and carbon compounds) and water exchange sites among the symbionts (Ivanov et al., 2019). Several AMF species, including Glomeraceae and Paraglomeraceae, produce vesicles the outer to middle cortical layers of root. Vesicles are storage structures contain lipids and glycogen and have a thin to thick wall layer (Smith & Read 2008). AMF structures in host plant roots are usually not observed without appropriate staining (Fig. 2).




Fig. 2. Structures of arbuscular mycorrhizal fungi inside and outside of host plant roots. a: Spores; b: Intra-radical hyphae (Hy) and Vesicle (Ve); c: Extra-radical hyphae (Ex-Hy); d: Hyphae (Hy) and Arbuscule (Ar) (Picture by the researcher)

Intra-radical and extra-radical hyphae of AMF are composed of aseptate multinuclear mycelia (Redecker et al., 2013). AMF spores are produced singly, in loose clusters, tight clusters, or sporocarps in the soil or within host plant roots (Schüßler & Walker 2010). Molecular studies have suggested that spores AMF are multinucleate, and a single

spore may contain multiple sequences of the same gene. Moreover, genetic variation of AMF is high not only between species but also nuclei in a single spore, and DNA polymorphism within fungal isolates was identified (Fitter 2005). To date, sexual reproduction has not been observed in AMF, although hidden sexual events were recently

suggested to happen (Corradi & Brachmann 2017). It was hypothesized the exchange of genetic material is carried out through anastomosis between AMF mycelia (Chagnon 2014, Novais et al., 2017).

History of the taxonomy of AMF

Fossil records and molecular evidence suggest that AMF symbiotic relationship with host plants formed 400–480 million years ago (between the Ordovician and the Devonian) and would have had essential roles in the establishment of terrestrial plants on land (Redecker et al., 2000). The symbiosis name of arbuscular mycorrhiza has changed over the years. First, the symbiosis was called “phycomycetous endomycorrhiza” to distinguish it from the endomycorrhizal symbioses, that formed between members of the Orchidaceae or Ericaceae and a fungus of endotrophic mycorrhiza. After discovering fungal structures, vesicles and arbuscules, the name “vesicular-arbuscular mycorrhiza” was established and persisted until recently. With regard to not all fungi forming vesicles, this symbiosis was renamed “arbuscular mycorrhiza” (Koide & Mosse 2004).

In recent years, the taxonomy and systematics of the AMF have experienced radical transformations. Kamienski (1881) discovered the symbiotic relationship between fungi and the roots of *Monotropa hypopitys*. Frank (1885) later used the term “mycorrhiza” for this relationship which means ‘fungus root.’ The Tulasne brothers (Tulasne & Tulasne 1845) were the first to describe the genus *Glomus* (*G. microcarpus* and *G. macrocarpus*) that considered this genus phylogenetically close to *Endogone*, a genus established by Link in 1809. Thaxter (1922) revised the Endogonaceae family, that included the genera *Endogone*, *Glaziella*, *Sclerocystis*, and *Sphaerocreas*. Peyronel (1923) was the first to suggest that the term “vesicular–arbuscular mycorrhizae” for members of the genus *Endogone*. Mosse (1953) reported the first study of adding the sporocarps of *Endogone* species to sterile soils with strawberry seedlings that resulted in mycorrhizal colonization.

Mosse (1953) reported the first study of inoculation the sporocarps of *Endogone* species to strawberry seedlings in sterile soil, which led to mycorrhizal colonization. This fungus was later named *Endogone mosseae* (= *Glomus mosseae* = *Funneliformis mosseae*) in her honor. After the device of the widely used wet sieving method by Gerdemann & Nicolson (1963) to extract fungal spores and sporocarps from soils, the interest in taxonomy of AMF increased dramatically. In the early 1970s, Gerdemann & Trappe split *Endogone* into seven nonmycorrhizal (*Endogone*, *Modicella*, *Glaziella*) and mycorrhizal (*Glomus*, *Sclerocystis*, *Gigaspora*, *Acaulospora*) genera, that were all placed in the family Endogonaceae, order Mucorales and phylum Zygomycota. As *Glaziella* and *Modicella* were not forming arbuscular mycorrhizal

symbiosis, Trappe (1982) transferred *Modicella* to the family Mortierellaceae, and Gibson et al. (1986) placed *Glaziella* in the Ascomycota. This classification provided a sound basis for a taxonomy of AMF for up to several years.

Ames & Schneider (1979) recognized and introduced a new genus, *Entrophospora*. Walker & Sanders (1986) separated a new genus called *Scutellospora* from the *Gigaspora*. For the ease of identification of AMF, Schenck and Perez (1988) have published “Manual for the Identification of VA Mycorrhizal Fungi”, which included all 126 AMF species descriptions. At the same time, different keys for the identification of AMF species developed. These manuals and keys are still being used in some laboratories to identify AMF species. During this time, AMF were identified based on morphological features and subcellular structures of spores. Morton & Benny (1990) placed the arbuscule-forming fungi into three families (Glomaceae, Acaulosporaceae, and Gigasporaceae) in a new order, the Glomales (now Glomerales). Cavalier-Smith (1998) proposed AMF could be placed in a new class, the Glomeromycetes within a new phylum Archeomycota. Analysis of available species of AMF and the examination of fossil records led to the suggestion of new taxa and the transfer of species to other genera. Taylor et al. (1995) proposed the genus *Glomites*. They described *Glomites rhyniensis* from aerial stems and rhizomes of the 400 million-year-old fossil plant *Aglaophyton major* in the Devonian period based on extra-radical and intraradical hyphae, chlamydospore-like spores, and arbuscule-like structures in the fossil plant.

Before molecular techniques, the only way to identify AMF was by microscopic examination of fungal spores. As early as 2001, Morton & Redecker erected two new families, Archaeosporaceae and Paraglomaceae in the order Glomales, by the data from molecular, morphological, and biochemical. Schüßler et al. (2001), using molecular data (SSU rDNA gene) and morphological characteristics, separated AMF from Zygomycota and grouped them in a monophyletic group, the phylum Glomeromycota, that is closely related to the Basidiomycota and the Ascomycota. Further, they proposed four new orders, i.e. Archaeosporales, Paraglomerales, Glomerales, and Diversisporales. Oehl & Sieverding (2004), identified four new species and after re-described two species of *Glomus*, classified them as a new genus *Pacispora* in the family Glomeraceae. In 2006, Sieverding & Oehl emended five species of *Entrophospora* and created the genera *Kuklospora* and *Intraspora*. Walker et al. (2007), based on morphological characteristics and rDNA data (SSU and ITS regions) proposed and described the genus *Ambispora* and the family Ambisporaceae. Oehl et al. (2008) introduced three new families in AMF

(Dentiscutataceae, Racocetraceae and, Scutellosporaceae).

Schüßler & Walker (2010) proposed a classification of AMF based on almost the entire length of the SSU rDNA gene. They separated *Glomus* into the genera *Funneliformis*, *Rhizophagus*, and *Sclerocystis* in the family Glomeraceae. They also erected the genus *Redeckera* in the family Diversisporaceae and the remaining species of *Glomus* along with *Claroideoglomus* in the family Claroideoglomeraceae. Oehl et al. (2011a) proposed a classification based on combined genetic (partial sequences of β -tubulin, and SSU and LSU rRNA) and phenotypic (traits associated with subtending hypha, e.g., color, shape and thickness, pore closure) characters. They divided the phylum Glomeromycota into three classes (Glomeromycetes, Archaeosporomycetes, and Paraglomeromycetes) and rearranged species of *Sclerocystis* and *Rhizophagus* and transferred back to the genus *Glomus*. In the same year, Oehl et al. (2011b) have been proposed new classes and orders. Goto et al. (2012), based on combined molecular and morphological studies, proposed a new classification and erected a new family called Intraornatosporaceae with the genera *Intraornatospora* and *Paradentiscutata*. Then, Kruger et al. (2012), based on morphological and molecular analyses of rDNA sequences (SSU, LSU, and ITS), divided this phylum into four orders: Paraglomales, Archaeosporales, Glomerales, and Diversisporales and ten genera. According to the classification made by Redecker et al. (2013), more than 250 species of AMF were reported, which were divided into one class (Glomeromycetes), four orders (Diversisporales, Glomerales, Archaeosporales, and Paraglomerales), 11 families and 25 genera. They rejected some genera like *Viscospora*, *Simiglomus*, and *Orbispora* and accepted *Dentiscutata*, *Sacculospora*, and *Corymbiglomus* but required verification. Blaszkowski et al. (2015) described two new genera, *Dominikia* and *Kamienskia*. Based on the phylogenetic analysis performed by Sieverding et al. (2015), *Rhizoglomus* genus forms a separate clade within the family Glomeraceae.

Taxonomy and classification of this fungal group have been rapidly updated over the last years, and several reports have been made. Walker et al. (2018) classified the phylum Glomeromycota into a single class (Glomeromycetes) which includes four orders (Diversisporales, Glomerales, Archaeosporales, and Paraglomerales), 12 families (Acaulosporaceae, Ambisporaceae, Archaeosporaceae, Claroideoglomeraceae, Diversisporaceae, Geosiphonaceae, Gigasporaceae, Glomeraceae, Pacisporaceae, Paraglomeraceae, Pervetustaceae, and Sacculosporaceae), 34 genera and approximately 316 species. In the same year, the

phylum Glomeromycota was divided by Tedersoo et al. into classes Glomeromycetes (Glomerales, Gigasporales, and Diversisporales), Paraglomeromycetes (Paraglomeromerales) and Archaeosporomycetes (Archaeosporales). Then, Wijayawardene et al. (2018) proposed another classification for Glomeromycota, which included two classes (Glomeromycetes, Archaeosporomycetes, and Paraglomeromycetes), four orders (Archaeosporales, Diversisporales, Glomerales, and Paraglomerales), 12 families and 38 genera and Gigasporales was not accepted. According to the latest unofficial classification presented on the website amf-phylogeny.com, AMF include four orders Diversisporales, Glomerales, Archaeosporales, and Paraglomerales, 12 families, 41 genera, and 342 species. Considering that new genera and species of this fungal group are being identified, such as *Microkamienskia* and *Microkamienskia peruviana* (Corazon-Guivin et al. 2019), the classification of Glomeromycota is debatable. Nevertheless, the classification performed by Wijayawardene et al. (2018) is used as a reference.

Morphological identification of AMF

Traditionally, AMF species have been identified by the morphological and anatomical characteristics of their spores (Traditionally, identification of AMF species has been made by spore extraction from soil and investigation of the morphological and anatomical characteristics of their spores) (Blaszkowski 2012). The morphological method is commonly used because it is fast, easy, cost-effective, and economical and can identify at the genus level. In the absence of spores, the intraradical and extra-radical structures of AMF (arbuscules, vesicles, hyphae, auxiliary cells) allow taxon identification at the family level (Merryweather & Fitter 1998, Sanders 2004).

There are several disadvantages to the morphological method: So far, an exact and comprehensive identification key has not been provided for AMF. AMF spores have simple structures, and morphological characteristics of the spores are scarce. For example, *Paraglomus* and *Glomus* cannot be distinguished from each other using spore morphology (Redecker et al., 2003). On the other hand, some species produce two types of spores or dimorphic spores (e.g., *Archaeospora leptoticha* and *G. dimorphicum*). Beside this, spore morphology may be high even within an AMF species (Merryweather & Fitter 1998). Also, spores collected from field soil may be degraded or parasitized and unidentifiable (Oehl et al., 2015, 2016). Another limitation of morphological identification is the spore production in this fungal group is strongly affected by environmental conditions and some species of AMF might sporulate rarely or only in a seasonal manner, such as *Acaulospora* spp. (Oehl et al., 2011b, 2012).

Aside from that, several AMFs may perform vegetative reproduction without the production of spores (Marleau et al., 2012). Thereby, the morphological diversity of AMF spores is insufficient for their variety in ecosystems. It is still common to the identification of AMF species by morphological features, notwithstanding the problems that are known in this method.

The first step in morphological identification of AMF species is the isolation of fungal. For this purpose, AMF spores were extracted from air-dried soil samples by wet sieving (Gerdemann & Nicolson 1963) and sucrose centrifugation (Jenkins 1964) methods. Intact and healthy spores are mounted on slides with PVLG + Melzer's reagent (1:1, v/v) (Hall 1984) and structural features are recorded, including spore color, shape, size, surface ornamentations, wall structure, hyphal attachments, and the mode of spore germination. Eventually, fungal spores are identified according to comprehensive identification manuals (Schenck & Perez 1988, Schüßler & Walker 2010, Oehl et al. 2011a, Wang & Liu 2017) and reference websites (<http://www.invam.wvu.edu>, <http://www.zor.zut.edu.pl>).

Molecular identification of AMF

In recent years, the development of new molecular methods has made it possible to identify symbiotic AMF species with different plant roots in a shorter period (Souza 2015). Molecular analysis is faster and provides more accurate and precise results. Molecular methods have revolutionized our understanding of phylogenetic relationships between fungi and fundamentally altered the morphological classification system (Hempel et al., 2007).

DNA extraction involves roots colonization, multiple spores, or single spore (Ferrol & Lanfranco 2020). The amount of AMF DNA extracted from different kinds of templates could lead to significant differences in the amplification of the target region (Shi et al., 2012). Field collected mycorrhizal roots contain various amounts of AMF tissue, and PCR inhibitors such as polysaccharides, proteins and phenolic compounds (Shafiqia & Stephan 2013). Also, the amount of fungal DNA compared to DNA plant is very low. So, selection of the suitable DNA extraction method and AMF-specific primers are necessary because otherwise, numerous saprophytic and pathogenic fungi will be identified simultaneously (Redecker et al. 2003).

Spores isolated from a soil sample are divided based on spore morphology using a stereomicroscope. The healthy spores that show numerous lipid globules inside, were selected and used for DNA extraction (Redecker 2020). Afterward, each spore type is grouped into three groups for molecular identification, morphological identification and purification. Cleaning spores by ultrasonic treatment and surface sterilization are helpful to remove debris and DNA from contaminating organisms (Sasvari et

al., 2012). DNA of spore (s) is extracted by crushing the spore using a micropipette. The most crucial reason for DNA extraction from a single spore, is reduced contaminations of other microbial species embedded in the spore wall. Removal of contamination attached to the single spore surface is more straightforward in comparison with multiple spores and mycorrhizal roots (Schwarzott et al., 2001).

Almost all molecular identification approaches for AMF are based on ribosomal DNA and the methods of Polymerase Chain Reaction (PCR) with specific primers (Kohout et al., 2014). Nested PCR using AMF-specific primers is a susceptible method that allows investigating of the mycorrhizal community structure and detection of fungal species present in roots and soil (Jacquot et al., 2000). Molecules other than rDNA, such as the mitochondrial genome, fatty acids, isozymes, and proteins (actin and β -tubulin), have been analyzed for AMF molecular identification (Raab 2007, Formey et al. 2012, Aminiannasab et al. 2021). Still, they are not widely used.

The ITS (internal transcribed spacer) region has been used for AMF molecular studies and sequencing (Opik et al., 2014, Tedersoo et al., 2014). But this region has a high degree of variation within AMF species, and it is difficult to find distinct features of closely related fungal species (Victorino et al., 2020). Therefore, ITS is of limited utility for AMF. Most data concerning molecular identification of AMF have been obtained using the SSU rDNA gene due to has a highly conserved gene (less variable than ITS) (Victorino et al. 2020) and provides intraspecific variation in this fungal group (Thiery et al. 2013, Lekberg et al. 2018, Perez-Lamarque et al. 2020). However, the slow evolution of this region is its most crucial disadvantage (Kruger et al., 2009, Bruns & Taylor 2016). Considering the LSU gene contains enough variation to distinguish species of AMF, this region has been chosen as a target for molecular identification and classification of AMF in some studies (House et al., 2016, Schutte et al., 2019). The use of the ITS variable region along with LSU and SSU regions, due to the high polymorphism, allows the identification of AMF to the species level and the differentiation of closely related species (Kruger et al. 2009). However, the SSU region is still the most frequent segment used in AMF molecular studies.

The selection of AMF-specific primers is essential as general primers will co-amplify the DNA of other organisms (Kruger et al., 2009). The first SSU sequences from rDNA of AMF were determined by Simon et al. (1992), who designed a specific primer for these fungi termed VANS1. The VANS1 primer was used in some studies (Clapp et al. 1995, Redecker et al. 2000), but it became clear that this primer is not well-conserved in all species of the AMF. Later, different specific primers were

designed to identify other species of AMF. Currently, the primer sets SSUmAf/LSUmAr and SSUmCf/LSUmBr designed by Kruger et al. (2009) are more specific and amplifies a partial on rDNA with a length of 1.5 kb. This 1.5 kb barcode consists of a part of the SSU, the complete ITS region (ITS1, 5.8S, ITS2 gene), and 800 bp of the LSU and avoids nonspecific amplification (Kruger et al., 2009). These primers should allow the distinction and separation of closely related AMF species such as *F. mosseae* and *F. coronatus* (Rosendahl & Matzen 2008), which the SSU or LSU regions alone cannot do (Schüßler 2013).

To evaluate the success of PCR amplification, PCR products are separated by electrophoresis through agarose gel in a TBE buffer. Much information about the sequences of rDNA of AMF species are accumulated in the GenBank database NCBI (<http://www.ncbi.nlm.nih.gov>). PCR-based techniques require continuous improvement in primer design as more sequences are introduced in the database. Each DNA sequence is compared with the nucleotide sequences in GenBank database NCBI using suitable software (Thompson et al., 1994). Sometimes the sequences obtained using PCR, may be unknown or uncultured Glomeromycota. This is because the rDNA sequences of some AMF species are not available in the GenBank database.

Today, critical progress has been made in the identification of AMF using morphological characteristics of AMF spores and molecular analysis targeting specific rDNA genes, which can easily be deduced from the numbers of AMF taxa detected (Njeru et al., 2015, Pontes et al., 2017). Molecular methods, due to their high identification accuracy can be complementary to morphological methods and lead to more confidence in identification and more stability in AMF taxonomy (Crossay et al., 2017).

Arbuscular Mycorrhizal Fungi in Iran

In Iran, many researchers have surveyed AMF communities and its diversity among different plant species. Since about 30 years ago, identification and description of AMF were made in Iran. For the first time, Mehravaran and Minasian (1984) identified 10 species of AMF, including *Rhizophagus fasciculatus* and *Gigaspora margarita*, from citrus orchards in the north and south of Iran. With the increase of information and the development of the methods used, several AMF species from different regions of Iran, combining molecular and morphological techniques, have been reported (e.g., Sedaghati et al., 2005; Rezaee Danesh et al., 2006; Sabet Jahromi et al., 2013; Yazdanpanah et al., 2017; Zangeneh 2021). However, most studies on AMF communities used morphological approaches and spores characteristics. Most AMF molecular identification systems used were based on rDNA genome and the Nested-PCR method. Among the

primers used can be mentioned to SSU-Glom1/LSU & ITS4/ITS5 (Salehi Jouzani et al. 2012; Sabet Jahromi et al. 2013), NS4/AML1 & AML1/AML2 (Yazdanpanah et al. 2017) and SSUmAf-LSUmAr/LSUmAr & SSUmCf/LSUmBr (Bazgir et al. 2020). It should be noted in the study conducted by Aminian et al. (2021) the partial sequences of the β -tubulin gene were amplified by three Nested-PCR using C2F/FBtub4R, GiH4R/IB36F and, Fsp/GiH3R primers.

Obviously, in Iran similar to other regions of the world, the composition and diversity of AMF species are affected by plant communities, soil and, climatic characteristics. These plants include food crops, vegetables, fruitful and non-fruitful trees, ornamental plants, medical plants, wild weeds, etc. In Table 1, some of the studies on identifying AMF species using morphological, molecular, and combination of both of these methods in Iran are listed. According to published articles, more than 115 AMF species have been identified at the species level (Table 2). Based on the results, the Glomeraceae species were dominant under different conditions, suggesting the higher and better adaptation of members of this family than other species. The predominance of Glomeraceae is consistent with the results of some studies in different countries (Ndoye et al., 2012, Bouazza et al., 2015, Mosbah et al., 2018). Members of some families, such as Acaulosporaceae, have occurred almost in the natural and intact areas. Moreover, it is probably the majority of AMF species that are still to be discovered and described.

CONCLUSION

The physiological and ecological benefits of mycorrhizal symbiosis for their host plants have long been acknowledged. The obligate biotrophic nature of AM fungi is one of the significant constraints of studying their taxonomy (Johnson et al., 2016). Detection and identification of AMFs based on spore morphology along with molecular methods provides a more comprehensive estimate of the composition, abundance, and diversity of the fungal community (Mosbah et al., 2018). Most AMF molecular identification studies have used primers that amplify ribosomal DNA, because the genes in this region have highly conserved coding sequences (Bati et al., 2015).

In Iran, many mycorrhizasts have done considerable amount of works on the identification of AMF species associated with plants located in different geographical regions of the country. They have used molecular and morphological methods to investigate the diversity and community composition of AMF symbiosis with varying plants in different areas. According to studies, the genera belonging to the family Glomeraceae, particular, *Glomus* and *Funneliformis*, are present in most areas of Iran that indicating greater adaptability of these genera to

varied soil conditions. In the future, AMF identification and classification must develop, and more applied research should be carried out to make

AMF commercially available in the agricultural sectors of Iran.

Table 1: AMF species/genera reported from different plants and regions in Iran

Plant/Region	Identification method	Result	Reference
Citrus	Morphological	Ten species	Mehravaran & Minassian 1984
Saffron	Morphological	15 species	Kianmehr et al., 1999
wheat, barley, maize, and sorghum	Morphological	21 species	Sadravi et al., 1999
Crops/Khorasan	Morphological	Six species	Balali et al., 2001
Grapes/Khorasan and Qazvin	Morphological	13 species	Sedaghati 2002
Different plants	Morphological	Five species	Sadravi 2002
Citrus	Morphological	Five species	Zangeneh et al., 2004
Kharturan	Morphological	12 species	Karimi et al., 2005
Sugarcane/Khuzestan	Morphological	17 species and two genera	Kariman et al., 2005
Pistachio/Rafsanjan	Morphological	Ten species and seven genera	Sedaghati et al., 2005
Citrus	Morphological	23 species	Zangeneh et al., 2005
Soybean	Morphological	21 species and five genera	Rezaee Danesh et al., 2006
Sugarcane	Morphological	14 species and four genera	Rokni et al., 2006
Wheat/Golestan	Morphological	19 species	Sadravi 2007
Sugarcane	Morphological	10 species and four genera	Rokni et al., 2010
Sugarcane	Morphological	16 species and four genera	Rokni & Goltapeh 2011
Grapevine/West Azerbaijan	Morphological	12 species and two genera	Mahdavi et al., 2012
Alfalfa/Kohgiluyeh & Boierahmad	Morphological	23 species and nine genera	Sadravi 2012
Wheat, barley and weeds	Morphological and Molecular	Seven species	Salehi Jouzani et al., 2012
Barley/Damghan	Morphological	16 species and four genera	Rezaee Danesh 2013
Wheat and barley	Morphological and Molecular	Nine species	Sabet Jahromi et al., 2013
<i>Crataegus pontica</i> / Ilam	Morphological	13 species and five genera	Mirzaei et al., 2014
Some Trees/Kiasar	Morphological	Ten species and three genera	Modarresi Chahardehi et al., 2014
Potato/Ardabil	Morphological	Four species and three genera	Rostami Hir et al., 2014
<i>Thymus daenensis</i>	Morphological	Nine species and two genera	Ahmadi et al., 2016
<i>Amygdalus scoparia</i>	Morphological	13 species and eight genera	Mirzaei & Moradi 2017
Pistachio/Kerman	Morphological and Molecular	Three species and three genera	Yazdanpanah et al., 2017
Manesht and Ghalarang protected area	Morphological	35 species and ten genera	Mirzaei et al., 2018
Pistachio/Rafsanjan	Morphological	Seven species and four genera	Aminizadeh et al., 2018
Saffron	Morphological and Molecular	Three species and three genera	Pooryousef et al., 2018
Medicinal plants/Kerman	Morphological and Molecular	Five species and four genera	Bazgir et al., 2020
Different plants/Rafsanjan	Morphological and Molecular	Five species and four genera	Aminiannasab et al., 2021
Different plants	Morphological	Five species and one genus	Zangeneh 2021

Table 2. List of number of the AMF species described in Iran

Families	Genus	Species
Glomeraceae	<i>Glomus</i>	<i>G. ambisporum</i> , <i>G. gibbosum</i> , <i>G. macrocarpum</i> , <i>G. pansihalos</i> , <i>G. sinuosum</i> , <i>G. microcarpum</i> , <i>G. badium</i> , <i>G. aureum</i> , <i>G. corymbiform</i> , <i>G. liquidambaris</i> , <i>G. botryoides</i> , <i>G. fulvum</i> , <i>G. gerdemanni</i> , <i>G. trimurales</i> , <i>G. albidum</i> , <i>G. rubiforme</i> , <i>G. fuegianum</i> , <i>G. dimorphicum</i> , <i>G. reticulatum</i> , <i>G. boreale</i> , <i>G. manihotis</i> , <i>G. tortuosum</i> , <i>G. invermaium</i> , <i>G. magnicaulis</i> , <i>G. pubescens</i> , <i>G. multisubstensum</i> ,
	<i>Septoglomus</i>	<i>Se. constrictum</i> , <i>Se. africanum</i> , <i>Se. deserticola</i> , <i>Se. turnauae</i> , <i>Se. xanthium</i> , <i>Se. jasnowskiae</i> , <i>Se. viscosum</i> , <i>Se. nakheelum</i>
	<i>Sclerocystis</i>	<i>Sc. coremioides</i>
	<i>Funneliformis</i>	<i>F. mosseae</i> , <i>F. caesaris</i> , <i>F. geosporus</i> , <i>F. caledonius</i> , <i>F. multiforus</i> , <i>F. coronatus</i> , <i>F. verruculosum</i> , <i>F. fragilistratum</i>
	<i>Rhizophagus</i>	<i>Rh. aggregatus</i> , <i>Rh. fasciculatus</i> , <i>Rh. intraradices</i> , <i>Rh. irregularis</i> , <i>Rh. diaphanus</i> , <i>Rh. clarus</i> , <i>Rh. microaggregatum</i> , <i>Rh. iranicus</i> , <i>Rh. arabicus</i>
	<i>Simiglomus</i>	<i>Si. hoi</i>
	<i>Dominikia</i>	<i>Do. disticha</i> , <i>Do. minuta</i>
	<i>Microkamienskia</i>	<i>M. perpusilla</i>
Claroideoglomeraceae	<i>Claroideoglomus</i>	<i>C. etunicatum</i> , <i>C. luteum</i> , <i>C. claroideum</i> , <i>C. walkeri</i> , <i>C. drummondii</i> , <i>C. lamellosum</i>
	<i>Viscospora</i>	<i>V. viscosa</i>
Diversisporaceae	<i>Diversispora</i>	<i>D. spurca</i> , <i>D. aurantia</i> , <i>D. eburnea</i> , <i>D. arenaria</i> , <i>D. varaderana</i> , <i>D. celata</i> , <i>D. pustulata</i> , <i>D. epigaea</i> , <i>D. versiformis</i> , <i>D. aurantium</i> , <i>D. globifera</i>
	<i>Desertispora</i>	<i>Des. omaniana</i>
Paraglomeraceae	<i>Paraglomus</i>	<i>Pa. laccatum</i> , <i>Pa. occultum</i> , <i>Pa. franciscana</i> , <i>Pa. scintillans</i>
Archeosporaceae	<i>Archaeospora</i>	<i>Ar. trappei</i> , <i>Ar. schenckii</i>
Acaulosporaceae	<i>Acaulospora</i>	<i>A. cavernata</i> , <i>A. mellea</i> , <i>A. dilatata</i> , <i>A. bireticulata</i> , <i>A. alpina</i> , <i>A. gedanensis</i> , <i>A. herrerae</i> , <i>A. punctata</i> , <i>A. scrobiculata</i> , <i>A. thomii</i> , <i>A. paulinae</i> , <i>A. capsicula</i> , <i>A. delicate</i> , <i>A. rugosa</i> , <i>A. colombiana</i>
Ambisporaceae	<i>Ambispora</i>	<i>Am. fennica</i>
Pacisporaceae	<i>Pacispora</i>	<i>P. boliviana</i> , <i>P. scintillans</i> , <i>P. franciscana</i>
Entrophosporaceae	<i>Entrophospora</i>	<i>E. infrequens</i>
Pervetustaceae	<i>Pervetustus</i>	<i>Pe. simplex</i>
Gigasporaceae	<i>Gigaspora</i>	<i>Gi. gigantea</i> , <i>Gi. rosea</i> , <i>Gi. decipiens</i> , <i>Gi. candida</i> , <i>Gi. albida</i> , <i>Gi. margarita</i>
	<i>Scutellospora</i>	<i>S. calospora</i> , <i>S. armeniaca</i> , <i>S. pellucida</i> , <i>S. coralloidea</i> , <i>S. fulgida</i> , <i>S. erythropha</i> , <i>S. dipurpurascens</i>
Dentiscutataceae	<i>Dentiscutata</i>	<i>De. reticulata</i> , <i>De. biornata</i>

REFERENCES

- Ahmadi T, Bernard F, Zangeneh S, Rejali, F. 2016. Identification of mycorrhizal fungi in the rhizosphere of *Thymus daenensis* and mycorrhization of this species with *Glomus intraradices* in green house conditions. Journal of Plant Research (Iranian Journal of Biology) 28(4): 674-683.
- Allen MF, Swenson W, Querejeta JJ, Egerton-Warburton LM, Treseder KK. 2003. Ecology of mycorrhizae: a conceptual framework for complex interactions among plants and fungi Annual Review of Phytopathology 41: 271-303.
- Al-Yahya'ei MN, Oehl F, Vallino M, Lumini E, Redecker D, Wiemken A, Bonfante P. 2011. Unique arbuscular mycorrhizal fungal communities uncovered in date palm plantations and surrounding desert habitats of Southern Arabia. Mycorrhiza 21: 195-209.
- Amballa H, Bhumi NR. 2016. Significance of arbuscular mycorrhizal fungi and rhizosphere microflora in plant growth and nutrition. In: Plant-microbe interaction: an approach to sustainable agriculture (pp. 417-452). Springer, Singapore.
- Aminiannasab P, Sedaghati E, Hosseini S, Saberi R. 2021. Identification of arbuscular mycorrhizal fungi associated with different plants in Rafsanjan based on morphological characteristics and β -tubulin gene sequence. Biological Journal of Microorganism 10(38): 27-40.
- Aminizadeh S, Hossein A, Ebrahim S, Mohammad M. 2018. Arbuscular mycorrhizal fungi associated with pistachio trees in Rafsanjan. Journal of Pistachio Science and Technology 1(2): 1-21.
- Ames RN, Schneider RW. 1979. *Entrophospora*, a new genus in the Endogonaceae. Mycotaxon 8: 347-352.
- Balali M, Kianmehr H, Mehravaran H, Hajian Shahri M. 2001. Introduction of 6 new species of mycorrhizal vesicular fungus (VAM) from Khorasan (Mashhad). Journal of Science Kharazmi University 1(2): 68-80.
- Balestrini R, Lumini E. 2018. Focus on mycorrhizal symbioses. Applied Soil Ecology 123: 299-304.
- Bati CB, Santilli E, Lombardo L. 2015. Effect of arbuscular mycorrhizal fungi on growth and on micronutrient and macronutrient uptake and allocation in olive plantlets growing under high total Mn levels. Mycorrhiza 25: 97-108.
- Bazgir E, Hatami N, Sedaghati E, Darvishnia M. 2020. Isolation and study of morphology and phylogeny of arbuscular mycorrhizal fungi coexisting with the roots of some medicinal plants in Kerman province. Agricultural Biotechnology Journal 12(1): 23-44.
- Begum N, Qin C, Ahanger MA, Raza S, Khan MI, Ashraf M, Ahmed N, Zhang L. 2019. Role of arbuscular mycorrhizal fungi in plant growth regulation: implications in abiotic stress tolerance. Frontiers in Plant Science 10: 1068-1085.
- Bencherif K, Boutekrabet A, Dalpe Y, Sahraoui AH. 2016. Soil and seasons affect arbuscular mycorrhizal fungi associated with Tamarix rhizosphere in arid and semi-arid steppes. Applied Soil Ecology 107: 182-190.
- Berruti A, Lumini E, Balestrini R, Bianciotto V. 2016. Arbuscular mycorrhizal fungi as natural biofertilizers: Let's benefit from past successes. Frontiers in Microbiology 6: 1-13.
- Binet MN, van Tuinen D, Deprêtre N, Koszela N, Chambon C, Gianinazzi S. 2011. Arbuscular mycorrhizal fungi associated with *Artemisia umbelliformis* Lam., an endangered aromatic species in Southern French Alps, influence plant P and essential oil contents. Mycorrhiza 21: 523-535.
- Bonfante P, Genre A. 2015. Arbuscular mycorrhizal dialogues: do you speak 'plantish' or 'fungish'? Trends in Plant Science 20(3): 150-154.
- Bouazza MK, Ighilhariz Z, de Lajudie P, Duponnois R, Bekki A. 2015. Assessing the native arbuscular mycorrhizal symbioses to rehabilitate a degraded coastal sand dune in Algeria. International Journal of Agriculture and Crop Sciences 8: 194-202.
- Blaszkowski J. 2012. Glomeromycota. W. Szafer Institute of Botany, Polish Academy of Sciences.
- Blaszkowski J, Chwat G, Goralska A, Ryszka P, Kovacs GM. 2015. Two new genera, *Dominikia* and *Kamienskia*, and *D. disticha* sp. nov. in Glomeromycota. Nova Hedwigia 100: 225-238.
- Cavalier-Smith T. 1998. A revised six-kingdom system of life. Biological Reviews 73: 203-266.
- Chagnon PL. 2014. Ecological and evolutionary implications of hyphal anastomosis in arbuscular mycorrhizal fungi. FEMS Microbiology Ecology 88(3): 437-444.
- Chen M, Arato M, Borghi L, Nouri E, Reinhardt D. 2018. Beneficial services of arbuscular mycorrhizal fungi—from ecology to application. Frontiers in Plant Science 9: 1270.
- Clapp JP, Young JP, Merryweather JW, Fitter AH. 1995. Diversity of fungal symbionts in arbuscular mycorrhizas from a natural community. New Phytologist 130: 259-265.
- Corazon-Guivin MA, Cerna-Mendoza A, Guerrero-Abad JC, Vallejos-Tapullima A, Carballar-Hernández S, da Silva GA, Oehl F. 2019. *Microkamienskia* gen. nov. and *Microkamienskia peruviana*, a new arbuscular mycorrhizal fungus from Western Amazonia. Nova Hedwigia 109(3-4): 335-368.
- Corradi N, Brachmann A. 2017. Fungal mating in the most widespread plant symbionts? Trends in Plant Science 22: 175-183.
- Crossay T, Antheaume C, Redecker D, Bon L, Chedri N, Richert C, Guentas L, Cavaloc Y, Amir H. 2017. New method for the identification of arbuscular mycorrhizal fungi by proteomic-based biotyping of spores using MALDI-TOF-MS. Scientific Reports 7(1): 1-16.
- Ferrol N, Azcon-Aguilar C, Perez-Tienda J. 2019. Arbuscular mycorrhizas as key players in sustainable plant phosphorus acquisition: An

- overview on the mechanisms involved Plant Science 280: 441-447.
- Ferrol N, Lanfranco L. 2020. Arbuscular mycorrhizal fungi: methods and protocols. Humana Press.
- Fitter AH. 2005. Darkness visible: reflections on underground ecology. Journal of Ecology 93: 231-43.
- Formey D, Moles M, Haouy A, Savelli B, Bouchez O, Becard G, Roux C. 2012. Comparative analysis of mitochondrial genomes of *Rhizophagus irregularis*-syn. *Glomus irregulare*-reveals a polymorphism induced by variability generating elements. New Phytologist 196: 1217-1227.
- Frank AB. 1885. Ueber die auf Wurzelsymbiose beruhende Ernahrung gewisser Baume durch unterirdische Pilze. Berichte der Deutschen Botanischen Gesellschaft 3: 128-145.
- Gao C, Montoya L, Xu L, Madera M, Hollingsworth J, Purdom E, Huttmacher RB, Dahlberg JA, Coleman-Derr D, Lemaux PG, Taylor JW. 2019. Strong succession in arbuscular mycorrhizal fungal communities. The ISME Journal 13: 214-226.
- Genre A, Lanfranco L, Perotto S, Bonfante P. 2020. Unique and common traits in mycorrhizal symbioses. Nature Reviews Microbiology 18: 649-660.
- Gerdemann J, Nicolson TH. 1963. Spores of mycorrhizal *Endogone* species extracted from soil by wet sieving and decanting. Transactions of the British Mycological Society 46: 235-244.
- Gerdemann JW, Trappe JM. 1974. The Endogonaceae in the Pacific Northwest. The New York Botanical Garden, New York.
- Gibson JL, Kimbrough JW, Benny GL. 1986. Ultrastructural observations on Endogonaceae (Zygomycetes). II. Glaziellales ord. nov. and Glaziellaceae fam. nov.: new taxa based upon light and electron microscopic observations of *Glaziella aurantiaca*. Mycologia 78: 941-954.
- Giovannini L, Sbrana C, Avio L, Turrini A. 2020. Diversity of a phosphate transporter gene among species and isolates of arbuscular mycorrhizal fungi. FEMS Microbiology Letters 367: 24.
- Gorzela MA, Holland TC, Xing X, Hart MM. 2012. Molecular approaches for AM fungal community ecology: a primer. Journal of Microbiological Methods 90: 108-114.
- Goto BT, Silva GA, Assis D, Silva DK, Souza RG, Ferreira AC, Jobim K, Mello C, Vieira HE, Maia LC, Oehl F. 2012. *Intraornatospora* (Gigasporales), a new family with two new genera and two new species. Mycotaxon 119: 117-132.
- Hall IR. 1984. Taxonomy of VA mycorrhizal fungi. CRC Press, Boca Raton, Florida, USA.
- Hayashi M, Niwa R, Urashima Y, Suga Y, Sato S, Hirakawa H, Yoshida S, Ezawa T, Karasawa T. 2018. Inoculum effect of arbuscular mycorrhizal fungi on soybeans grown in long-term bare-fallowed field with low phosphate availability. Soil Science and Plant Nutrition 64: 306-311.
- Hempel S, Renker C, Buscot F. 2007. Differences in the species composition of arbuscular mycorrhizal fungi in spore, root and soil communities in a grassland ecosystem. Environmental Microbiology 9: 1930-1938.
- House GL, Ekanayake S, Ruan Y, Schutte UM, Kaonongbua W, Fox G, Ye Y, Bever JD. 2016. Phylogenetically structured differences in rRNA gene sequence variation among species of arbuscular mycorrhizal fungi and their implications for sequence clustering. Applied and Environmental Microbiology 82: 4921-4930.
- Ivanov S, Austin J, Berg RH, Harrison MJ. 2019. Extensive membrane systems at the host-arbuscular mycorrhizal fungus interface. Nature Plants 5: 194-203.
- Jacquot E, Tuinen D van, Gianinazzi S, Gianinazzi-Pearson V. 2000. Monitoring species of arbuscular mycorrhizal fungi in planta and in soil by nested PCR: application to the study of the impact of sewage sludge. Plant and Soil 226(2): 179-188.
- Jenkins WR. 1964. A rapid centrifugal-flotation technique for separating nematodes from soil. Plant Disease Reporter 48(9): 692.
- Johnson JM, Houngnandan P, Kane A, Chatagnier O, Sanon KB, Neyra M, van Tuinen D. 2016. Colonization and molecular diversity of arbuscular mycorrhizal fungi associated with the rhizosphere of cowpea (*Vigna unguiculata* (L.) Walp.) in Benin (West Africa): an exploratory study. Annals of Microbiology 66: 207-221.
- Kamienski F. 1881. Die Vegetationsorgane der *Monotropa hypopitys* L. Botanische Zeitung 29: 457-461.
- Kariman KH, Goltapeh EM, Minassian V. 2005. Arbuscular mycorrhizal fungi from Iran. Journal of Agricultural Technology 1(2): 301-313.
- Karimi F, Zangeneh S, Yousefzadi M, Zarre Mayvan H. 2005. Recognition of arbuscular-mycorrhiza fungi (AMF) and root colonization percentage in Kharturan biosphere reserve. Environmental Sciences 10: 83-88.
- Kianmehr H, Gholami Moghadam F, Hajian Shahri M, Falahati Rastegar M. 1999. Report of new species of vesicular-arbuscular mycorrhiza fungi from Khorasan (northeastern Iran). Desert Journal 4(2): 25-44.
- Kohout P, Sudova R, Janouskova M, Ctvrtlikova M, Hejda M, Pankova H, Slavikova R, Stajerova K, Vosatka M, Sykorova Z. 2014. Comparison of commonly used primer sets for evaluating arbuscular mycorrhizal fungal communities: is there a universal solution? Soil Biology and Biochemistry 68: 482-493.
- Koide RT, Mosse B. 2004. A history of research on arbuscular mycorrhiza. Mycorrhiza 14(3): 145-163.
- Kruger M, Stockinger H, Kruger C, Schubler A. 2009. DNA-based species level detection of Glomeromycota: one PCR primer set for all

- arbuscular mycorrhizal fungi. *New Phytologist* 183: 212-223.
- Kruger M, Kruger C, Walker C, Stockinger H, Schussler A. 2012. Phylogenetic reference data for systematics and phylotaxonomy of arbuscular mycorrhizal fungi from phylum to species level. *New Phytologist* 193: 970-984.
- Lakmali JPD, Yapa PN, Sadaruwan MKKD. 2019. Role of the vetiver (*Chrysopogon zizanioides* L.) and arbuscular mycorrhizal fungi on soil quality improvement in *Pinus* plantation soil. In proceedings of International Research Symposium on Pure and Applied Sciences, Faculty of Science, University of Kelaniya, Sri Lanka. 25th October 2019.
- Lekberg Y, Vasar M, Bullington LS, Sepp SK, Antunes PM, Bunn R, Larkin BG, Opik M. 2018. More bang for the buck? Can arbuscular mycorrhizal fungal communities be characterized adequately alongside other fungi using general fungal primers? *New Phytologist* 220(4): 971-976.
- Lovelock CE, Andersen K, Morton JB. 2003. Arbuscular mycorrhizal communities in tropical forests are affected by host tree species and environment. *Oecologia* 135: 268-279.
- Li J, Awasthi MK, Xing W, Liu R, Bao H, Wang X, Wang J, Wu F. 2020. Arbuscular mycorrhizal fungi increase the bioavailability and wheat (*Triticum aestivum* L.) uptake of selenium in soil. *Industrial Crops and Products* 150: 112383.
- Mahdavi A, Rezaee Danesh Y, Abbasi A, Mahdavi F, Jamaati-e-Somarin S. 2012. Study on occurrence and diversity of Arbuscular mycorrhizal fungi associated with grapevine rhizosphere in West Azerbaijan Province in Iran. *African Journal of Microbiology Research* 6(6): 1238-1242.
- Marleau J, Dalpe Y, St-Arnaud M, Hijri M. 2011. Spore development and nuclear inheritance in arbuscular mycorrhizal fungi. *BMC Evolutionary Biology*, 11(1): 1-11.
- Mehravaran H, Minassian H. 1984. Study of mycorrhizal citrus in Iran. Proceedings of the 9th Iranian Congress of Medicinal Plants. University of Tehran, Tehran, Iran.
- Merryweather J, Fitter A. 1998. The arbuscular mycorrhizal fungi of *Hyacinthoides non-scripta* I. Diversity of fungal taxa. *New Phytologist* 138: 117-129.
- Mirzaei J, Noorbakhsh N, Karamshahi A. 2014. Identification of Arbuscular Mycorrhizal Fungi Associated with *Crataegus pontica* C. Koch from Ilam Province, Iran. *ECOPERSIA* 2(4): 767-777.
- Mirzaei J, Moradi M. 2017. Biodiversity of arbuscular mycorrhizal fungi in *Amygdalus scoparia* Spach plantations and a natural stand. *Journal of Forestry Research* 28(6): 1209-1217.
- Mirzaei J, Dostcami S, Moradi M. 2018. Identification of arbuscular mycorrhizal fungi associated with plant species in the Manesht and Ghalarang protected area. *Forest and Wood Products* 70(4): 549-557.
- Modarresi Chahardehi A, Mousavi L, Bakhtkhan Ardeh Jani T, Rezaee Danesh Y, Ibrahim D. 2014. Isolation and identification of dominant arbuscular mycorrhizal fungi in some trees in region of Kiasar. *Iranian Journal of Plant Protection Science* 45(2): 371-380.
- Morton JB, Benny GL. 1990. Revised classification of arbuscular mycorrhizal fungi (Zygomycetes): a new order, Glomales, two new suborders, Glomineae and Gigasporineae, and two new families, Acaulosporaceae and Gigasporaceae, with an emendation of Glomeraceae. *Mycotaxon* 37: 471-491.
- Morton JB, Redecker D. 2001. Two families of Glomales, Archaeosporaceae and Paraglomeraceae, with two new genera *Archaeospora* and *Paraglomus*, based on concordant molecular and morphological characters. *Mycologia* 93: 181-195.
- Mosbah M, Philippe DL, Mohamed M. 2018. Molecular identification of arbuscular mycorrhizal fungal spores associated to the rhizosphere of *Retama raetam* in Tunisia. *Soil Science and Plant Nutrition* 64(3): 335-341.
- Mosse B. 1953. Fructifications associated with mycorrhizal strawberry roots. *Nature* 171: 974.
- Ndoye F, Kane A, Ngonkeu Mangaptche EL, Bakhoum N, Sanon A, Diouf D, Sy MO, Baudoin E, Noba K, Prin Y. 2012. Changes in land use system and environmental factors affect arbuscular mycorrhizal fungal density and diversity, and enzyme activities in rhizospheric soils of *Acacia senegal* (L.) Willd. *International Scholarly Research Notices* 2012: 1-14.
- Nielsen KB, Kjoller R, Bruun HH, Schnoor TK, Rosendahl S. 2016. Colonization of new land by arbuscular mycorrhizal fungi. *Fungal Ecology* 20: 22-29.
- Njeru E, Avio L, Bocci G, Sbrana C, Turrini A, Barberi P, Giovannetti M, Oehl F. 2015. Contrasting effects of cover crops on 'hot spot' arbuscular mycorrhizal fungal communities in organic tomato. *Biology and Fertility of Soils* 51: 151-166.
- Novais CBD, Pepe A, Siqueira JO, Giovannetti M, Sbrana C. 2017. Compatibility and incompatibility in hyphal anastomosis of arbuscular mycorrhizal fungi. *Scientia Agricola* 74: 411-416.
- Oehl F, Sieverding E. 2004. *Pacispora*, a new vesicular arbuscular mycorrhizal fungal genus in the Glomeromycetes. *Journal of Applied Botany* 78: 72-82.
- Oehl F, de Souza FA, Sieverding E. 2008. Revision of *Scutellospora* and description of five new genera and three new families in the arbuscular mycorrhiza forming Glomeromycetes. *Mycotaxon* 106: 311-360.
- Oehl F, Alves a Silva G, Goto BT, Costa Maia L, Sieverding E. 2011a. Glomeromycota: two new

- classes and a new order. *Mycotaxon* 116(1): 365-379.
- Oehl F, Schneider D, Sieverding E, Burga CA. 2011b. Succession of arbuscular mycorrhizal communities in the foreland of the retreating Morteratsch glacier in the Central Alps. *Pedobiologia* 54: 321-331.
- Oehl F, Palenzuela J, Sanchez-Castro I, Kuss P, Sieverding E, Silva GA. 2012. *Acaulospora nivalis*, a new fungus in the Glomeromycetes, characteristic for high alpine and nival altitudes of the Swiss Alps. *Nova Hedwigia* 95: 105-122.
- Oehl F, Sanchez-Castro I, Palenzuela J, Silva GA. 2015. *Palaeospora spainii*, a new arbuscular mycorrhizal fungus from Swiss agricultural soils. *Nova Hedwigia* 101: 89-102.
- Oehl F, Santos VM, Palenzuela J. 2016. *Paraglomus turpe*, a new arbuscular mycorrhizal fungal species from Central European agricultural soils. *Nova Hedwigia* 103: 491-499.
- Opik M, Davison J, Moora M, Zobel M. 2014. DNA-based detection and identification of Glomeromycota: the virtual taxonomy of environmental sequences. *Botany-Botanique* 92: 135-147.
- Perez-Lamarque B, Opik M, Maliet O, Silva ACA, Selosse M, Martos F, Morlon H. 2020. Global drivers of obligate mycorrhizal symbionts diversification. *bioRxiv* 2020.
- Peyronel B. 1923. Fructification de l'endophyte a arbuscules et a vesicules des mycorrhizes endotrophes. *Bulletin Society Mycological France* 39: 119-126.
- Pontes JS, Oehl F, Marinho F, Coyne D, Silva DKA, Yano-Melo AM, Maia LC. 2017. Diversity of arbuscular mycorrhizal fungi in Brazil's Caatinga and experimental agroecosystems. *Biotropica* 49: 413-427.
- Pooryousef H, Hemmat J, Vaez M. 2018. A comparison between molecular and morphological methods in evaluation of vesicular-arbuscular mycorrhizal fungi colonization with *Crocus sativus* L (saffron). *Biological Journal of Microorganism* 7(26): 1-10.
- Raab PA. 2007. Development of new molecular markers for phylogeny and molecular identification of arbuscular mycorrhizal fungi (Glomeromycota). PhD thesis, University of Basel.
- Redecker D, Kodner R, Graham LE. 2000. Glomalean fungi from the Ordovician. *Science* 289:1920-1921.
- Redecker D, Hijri I, Wiemken A. 2003. Molecular identification of arbuscular mycorrhizal fungi in roots: perspectives and problems. *Folia Geobotanica* 38: 113-124.
- Redecker D, Schüßler A, Stockinger H, Sturmer SL, Morton JB, Walker C. 2013. An evidence-based consensus for the classification of arbuscular mycorrhizal fungi (Glomeromycota). *Mycorrhiza* 23(7): 515-531.
- Redecker D. 2020. Single-spore extraction for genetic analyses of arbuscular mycorrhizal fungi. In: *Arbuscular mycorrhizal fungi. methods in molecular biology*. (N Ferrol, L Lanfranco, eds). Humana, New York.
- Rezaee Danesh Y, Goltapeh EM, Alizadeh A, Varma A, Mukerjee KG. 2006. Distribution and abundance of arbuscular mycorrhizal fungi from soybean rhizosphere in Iran. *Journal of Agricultural Technology* 2: 251-257.
- Rezaee Danesh Y. 2013. Evaluation of arbuscular mycorrhizal fungi with barley in Damghan region. *Journal of Plant Protection* 26(4): 437-449.
- Rokni N, Goltapeh EM, Alizadeh A. 2006. Introducing one species of arbuscular mycorrhizal fungus from Khuzestan sugarcane fields new to Iran. *Rostaniha* 7(2): 15-21.
- Rokni N, Goltapeh EM, Alizadeh A. 2010. Some new recorded species of arbuscular mycorrhizal fungi associated with sugarcane crop in Iran. *Journal of Agricultural Technology* 6: 67-78.
- Rokni N, Goltapeh EM. 2011. Diversity of arbuscular mycorrhizal fungi associated with common sugarcane varieties in Iran. *Journal of Agricultural Technology* 7: 1017-1022.
- Rosendahl S, Matzen HB. 2008. Genetic structure of arbuscular mycorrhizal populations in fallow and cultivated soils. *New Phytologist* 179: 1154-1161.
- Rostami Hir S, Riahi H, Zanganeh S. 2014. Identification of mycorrhiza and effect on the growth parameters potato. *Environmental Sciences* 12(4): 75-80.
- Sabet Jahromi M, Salehi Jouzani Gh, Hosseini M, Khayam Nekoei SM, Akbari Vala S. 2013. Isolation and identification of native mycorrhizal fungi in agricultural soils in arid regions of Iran using nested PCR. *Journal of Biotechnology, Tarbiat Modarres University* 1: 1-13.
- Sadravi M. 1999. Identification of arbuscular mycorrhizal fungi of wheat, barley, maize and sorghum in Tehran and Khuzestan provinces. M.Sc. dissertation, Tarbiat Modares University, Tehran, Iran.
- Sadravi M. 2002. Symbiotic fungi of soybean root, sunflower and sesame in Golestan province. The first national conference and festival of oilseeds.
- Sadravi M. 2007. Arbuscular mycorrhizal fungi in wheat fields in Golestan province. *Vegetables* 7: 129-140.
- Sadravi M. 2012. Arbuscular mycorrhizal fungi of alfalfa root in the Kohgiluyeh and Boierahmad province (SW Iran). *Rostaniha* 13(1): 101-104.
- Salehi Jouzani Gh, Akbari Vala S, Sabet Jahromi M, Morsali H. 2012. Isolation and identification of dominant arbuscular mycorrhizal fungi in the rhizosphere of wheat, barley and weeds in some saline regions of Iran. *Journal of Crop Biotechnology* 1(1): 61-75.
- Sanders IR. 2004. Plant and arbuscular mycorrhizal fungal diversity are we looking at the relevant levels of diversity and are we using the right techniques? *New Phytologist* 164: 415-418

- Sasvari Z, Magurno F, Galanics D, Hang TT, Ha TT, Luyen ND, Posta K. 2012. Isolation and identification of arbuscular mycorrhizal fungi from agricultural fields of Vietnam. *American Journal of Plant Sciences* 3: 1796-1801.
- Sieverding E, Oehl F. 2006. Revision of *Entrophospora* and description of *Kuklospora* and *Intraspora*, two new genera in the arbuscular mycorrhizal Glomeromycetes. *Journal of Applied Botany and Food Quality* 80: 69-81.
- Schenck NC, Perez Y. 1988. Manual for the identification of VA mycorrhizal fungi. Plant Pathology Department Institute of Food and Agricultural Sciences University of Florida.
- Schüßler A, Schwarzott D, Walker C. 2001. A new fungal phylum, the Glomeromycota phylogeny and evolution. *Mycological Research* 105: 1413-1421.
- Schüßler A, Walker C. 2010. The Glomeromycota: a species list with new families and new genera. The Royal Botanic Garden, Edinburgh.
- Schüßler A. 2013. Phylogeny and taxonomy of Glomeromycota.
- Schutte UM, Henning JA, Ye Y, Bowling A, Ford J, Genet H, Waldrop MP, Turetsky MR, White JR, Bever JD. 2019. Effect of permafrost thaw on plant and soil fungal community in a boreal forest: does fungal community change mediate plant productivity response? *Journal of Ecology* 107: 1737-1752.
- Schwarzott D, Walker C, Schussler A. 2001. *Glomus*, the largest genus of the arbuscular mycorrhizal fungi (Glomales), is nonmonophyletic. *Molecular Phylogenetics and Evolution* 21: 190-197.
- Sedaghati E. 2002. Isolation and identification of arbuscular mycorrhizal fungi in the Khorasan and Qazvin provinces. M.Sc. dissertation, Tarbiat Modares University, Tehran, Iran.
- Sedaghati E, Hosseini A, Khodayegan P, Raghani M, Maohammadi E. 2005. Report of some species of arbuscular Mycorrhiza fungi from Pistachio orchards Rafsanjan, Iran. International Symposium on Pistachio and Almond-ISHS-Tehran-Iran. 202.
- Shafiqia G, Stephan R. 2013. Molecular detection of Arbuscular mycorrhizal fungi in calcareous soil and roots of *Zea mays* L. *Journal of Academia and Industrial Research (JAIR)* 1(19): 547-549.
- Shi P, Abbott LK, Banning NC, Zhao B. 2012. Comparison of morphological and molecular genetic quantification of relative abundance of arbuscular mycorrhizal fungi within roots. *Mycorrhiza* 22(7): 501-513.
- Simon L, Levesque RC, Lalonde M. 1992. Identification of endomycorrhizal fungi colonizing roots by fluorescent single-strand conformation polymorphism-polymerase chain reaction. *Applied and Environmental Microbiology* 59: 4211-4215.
- Sieverding E, da Silva GA, Berndt R, Oehl F. 2015. *Rhizoglomus*, a new genus of the Glomeraceae. *Mycotaxon* 129(2): 373-386.
- Smith SE, Read DJ. 2008. Mycorrhizal symbiosis. UK. London, Academic Press, 125 pp.
- Smith SE, Smith FA. 2011. Roles of arbuscular mycorrhizas in plant nutrition and growth: new paradigms from cellular to ecosystem scales. *Annual Review of Plant Biology* 62: 227-250.
- Souza T. 2015. Handbook of arbuscular mycorrhizal fungi (1st ed). Springer International Publishing, Switzerland 4: 87-128.
- Symanczik S. 2016. Arbuscular mycorrhizal fungal diversity of arid lands: from Am fungal species to AM fungal communities PhD thesis, Basel University, Switzerland.
- Tahat MM, Sijam K. 2012. Arbuscular mycorrhizal fungi and plant root exudates bio-communications in the rhizosphere. *African Journal of Microbiology Research* 6(46): 7295-7301.
- Tao L, Ahmad A, De Roode JC, Hunter MD. 2016. Arbuscular mycorrhizal fungi affect plant tolerance and chemical defences to herbivory through different mechanisms. *Journal of Ecology* 104: 561-571.
- Taylor TN, Remy W, Hass H, Kerp, H. 1995. Fossil arbuscular mycorrhizae from the Early Devonian. *Mycologia* 87: 560-573.
- Tedersoo L, Bahram M, Polme S, Koljalg U, Yorou NS, Wijesundera R, Ruiz LV, Vasco-Palacios AM, Thu PQ, Suija A, Smith ME. 2014. Global diversity and geography of soil fungi. *Science* 346: 1256688.
- Tedersoo L, Sanchez-Ramirez S, Koljalg U, Bahram M, Doring M, Schigel D, May T, Ryberg M, Abarenkov K. 2018. High-level classification of the Fungi and a tool for evolutionary ecological analyses. *Fungal Diversity* 90(1): 135-159.
- Thaxter R. 1922. A revision of the *Endogoneae*. *Proceeding of American Academy of Arts and Sciences* 57(12): 291-350.
- Thompson JD, Higgins DG, Gibson TJ. 1994. CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Research* 22: 4673-4680.
- Trappe JM. 1982. Synoptic keys to the genera and species of zygomycetous mycorrhizal fungi. *Phytopathology* 72(8): 1102-1108.
- Tulasne LR, Tulasne C. 1844. Fungi nonnulli hipogaei, novl v. minus cogniti auct. *Giornale Botanico Italiano* 2(1): 55-63.
- Turrini A, Agnolucci M, Palla M, Tomé E, Tagliavini M, Scandellari F, Giovannetti M. 2017. Species diversity and community composition of native arbuscular mycorrhizal fungi in apple roots are affected by site and orchard management. *Applied Soil Ecology* 116: 42-54.
- Varga S, Finozzi C, Vestberg M, Kytoviita MM. 2015. Arctic arbuscular mycorrhizal spore community and viability after storage in cold conditions. *Mycorrhiza* 25(5): 335-343.

- Victorino IMM, Berruti A, Orgiazzi A, Voyron S, Bianciotto V, Lumini E. 2020. High-Throughput DNA Sequence-Based Analysis of AMF Communities. In: Arbuscular mycorrhizal fungi. methods in molecular biology. (N Ferrol, L Lanfranco, eds). Humana, New York.
- Walker CH, Sanders FE. 1986. Taxonomic concepts in the Endogonaceae. III. The separation of *Scutellosopra* gen. nov. from *Gigaspora* Gerd. and Trappe. *Mycotaxon* 27: 169-182.
- Walker C, Vestberg M, Demircik F, Stockinger H, Saito M, Sawaki H, Nishimura I, Schüßler A. 2007. Molecular phylogeny and new taxa in the Archaeosporales (Glomeromycota): *Ambispora fennica* gen. sp. nov., *Ambisporaceae* fam. nov., and emendation of *Archaeospora* and *Archaeosporaceae*. *Mycological Research* 111: 137-153.
- Walker C, Gollotte A, Redecker D. 2018. A new genus, *Planticonsortium* (Mucoromycotina), and new combination (*P. tenue*), for the fine root endophyte, *Glomus tenue* (basionym *Rhizophagus tenuis*). *Mycorrhiza* 28: 213-219.
- Wang Y, Liu R. 2017. A checklist of arbuscular mycorrhizal fungi in the recent taxonomic system of Glomeromycota. *Mycosystema* 36(7): 820-850.
- Wang J, Ma S, Wang GG, Xu L, Fu Z, Song J, Zhang J. 2021. Arbuscular mycorrhizal fungi communities associated with wild plants in a coastal ecosystem. *Journal of Forestry Research* 32(2): 683-695.
- Wetzel K, Silva GA, Matczinski U, Oehl F, Fester T. 2014. Superior differentiation of arbuscular mycorrhizal fungal communities from till and no-till plots by morphological spore identification when compared to T-RFLP. *Soil Biology and Biochemistry* 72: 88-96.
- Wijayawardene NN, Pawłowska J, Letcher PM, Kirk PM, Humber RA, Schüßler A, Wrzosek M, Muszewska A, Okrasinska A, Istel L, Gęsiorska A. 2018. Notes for genera: basal clades of Fungi (including Aphelidiomycota, Basidiobolomycota, Blastocladiomycota, Calcarisporiellomycota, Caulochytriomycota, Chytridiomycota, Entomophthoromycota, Glomeromycota, Kickxellomycota, Monoblepharomycota, Mortierellomycota, Mucoromycota, Neocallimastigomycota, Olpidiomycota, Rozellomycota and Zoopagomycota). *Fungal Diversity* 92(1): 43-129.
- Willis A, Rodrigues BF, Harris PJ. 2013. The ecology of arbuscular mycorrhizal fungi. *Critical Reviews in Plant Sciences* 32(1): 1-20.
- Wu MJ, Yan YR, Wang YQ, Mao Q, Fu YL, Peng XH, Yang ZK, Ren JJ, Liu AR, Chen SC, Ahammed GJ. 2021. Arbuscular mycorrhizal fungi for vegetable (VT) enhance resistance to *Rhizoctonia solani* in watermelon by alleviating oxidative stress. *Biological Control* 152: 104433.
- Yazdanpanah M, Sedaghati E, Khodygan P, Alaei H, Nadi M. 2017. Molecular and morphological identification of arbuscular mycorrhizal fungi associated with pistachio roots in Kerman province. *Journal of Pistachio Science and Technology* 2(4): 11-29.
- Zangeneh S, Shirvani A, Alian Y. 2004. Introducing new species of arbuscular-mycorrhizal fungi from the Iranian citrus rhizosphere. *Vegetables* 6: 8-77.
- Zangeneh S, Shirvani A, Alian Y, Najafi Nia M, Karam Pur F, Ghale Dozdani H. 2005. Introduction of some new arbuscular-mycorrhizal fungi (AMF) from citrus rhizosphere of Iran. *Rostaniha* 6: 8-77.
- Zangeneh S. 2021. Introduction of some new species of genus *Acaulospora* from Iran. *Rostaniha* 22(1): 67-74.
- Zhang F, Liu M, Li Y, Chen YY, Xiao Y. 2019. Effects of arbuscular mycorrhizal fungi, biochar and cadmium on the yield and element uptake of *Medicago sativa*. *Science of The Total Environment* 655: 1150-1158.
- Zhang F, Zou YN, Wu QS, Kuca K. 2020. Arbuscular mycorrhizas modulate root polyamine metabolism to enhance drought tolerance of trifoliate orange. *Environmental and Experimental Botany* 171: 103962.
- Zou YN, Wu QS, Kuca K. 2021. Unravelling the role of arbuscular mycorrhizal fungi in mitigating the oxidative burst of plants under drought stress. *Plant Biology* 23: 50-57.

مروری بر مطالعات تاکسونومیکی قارچ‌های میکوریز آربوسکولار در ایران

ابراهیم صداقتی^۱، مرضیه یزدان‌پناه^۲، ماریه نادی^۳

۱. گروه بیماری‌شناسی، دانشکده کشاورزی، دانشگاه ولی عصر (عج) رفسنجان، ایران

۲. گروه بیماری‌شناسی گیاهی، دانشگاه علوم کشاورزی و منابع طبیعی گرگان، ایران

۳. مرکز تحقیقات پسته، پژوهشکده علوم باغبانی، سازمان آموزش و ترویج تحقیقات کشاورزی، رفسنجان، ایران

چکیده: قارچ‌های میکوریز آربوسکولار (AMF) یکی از مهم‌ترین ریزجانداران مفید خاک هستند که قادر به برقراری رابطه همزیستی همیاری با طیف وسیعی از گیاهان می‌باشند. AMF کربن مورد نیاز برای تکمیل چرخه زندگی خود را از گیاه میزبان دریافت می‌کنند. در مقابل، این گروه قارچی فواید متعددی برای گیاهان میزبان دارند، از جمله محافظت از گیاه در برابر بیمارگرها، افزایش تحمل تنش‌های غیرزیستی (خشکی و شوری) و افزایش جذب آب و مواد مغذی. در گذشته، شناسایی و طبقه‌بندی AMF عمدتاً براساس ویژگی‌های ریخت‌شناسی اسپورها مانند نحوه تشکیل، ساختار دیواره و ویژگی‌های هیف اتصال انجام می‌گردید. سپس، روش‌های مولکولی برای شناسایی این قارچ‌ها مورد استفاده قرار گرفتند. روش‌های مبتنی بر PCR منجر به شناسایی مستقیم گونه‌های AMF موجود در ریشه‌های گیاه یا ناحیه فراریشه می‌شوند. تاکنون، پرایمرهای متعددی برای افزایش دقت شناسایی AMF طراحی شده است. امروزه، طبقه‌بندی این گروه قارچی مبتنی بر روش‌های ریخت‌شناسی و مولکولی می‌باشد. در این مطالعه، مروری بر پژوهش‌های انجام شده در زمینه شناسایی AMF در ایران ارائه می‌گردد. طی چند دهه اخیر، شناسایی AMF در ایران مورد توجه فزاینده‌ای قرار گرفته است. بدین منظور، پژوهش‌گران از ویژگی‌های ریخت‌شناسی و روش‌های مبتنی بر DNA برای شناسایی این گروه قارچی استفاده کرده‌اند. تاکنون بیش از ۱۱۵ گونه AMF متعلق به ۲۲ جنس از مناطق و جوامع گیاهی مختلف مانند زمین‌های زراعی، جنگل‌ها، مراتع و غیره، شناسایی و گزارش شده است. شناخت ترکیب و تنوع AMF برای استفاده از آن‌ها به عنوان کودهای زیستی در کشاورزی جهت کاهش نهاده‌های شیمیایی و افزایش تولید محصول حائز اهمیت می‌باشد.

کلمات کلیدی: ترکیب جامعه، تنوع، ریخت‌شناسی اسپور، همزیستی، *Glomus*.