



Bioactive potential and GC-MS fingerprinting of extracts from endophytic fungi associated with seeds of some medicinal plants

A. Asghari

M.A. Tajick Ghanbary ✉

Department of Plant Protection, Faculty of Crop Sciences, Sari Agricultural Sciences and Natural Resources University, Sari, Iran.

M. Bakhshi

Department of Botany, Iranian Research Institute of Plant Protection, Agricultural Research, Education and Extension Organization (AREEO), Tehran, Iran.

V. Babaeizad

Department of Plant Protection, Faculty of Crop Sciences, Sari Agricultural Sciences and Natural Resources University, Sari, Iran.

Abstract: Endophytic fungi are a group of host-associated fungal communities that benefit their hosts. According to the conditions of their specific living environment, plant endophytic fungi produce many bioactive metabolites with different structural features. The bioactive compounds isolated from endophytic fungi, have significant effects on increasing the compatibility of both endophytic fungi and their host plants, such as the tolerance to biotic and abiotic stresses. In addition, some of these metabolites have indicated medicinal and ecological importance. In the present investigation, five fungal endophytes were isolated from the seeds of some medicinal plants. These endophytic isolates were characterized by sequencing of Internal Transcribed Spacer (ITS) regions as *Acremonium* sp. isolated from *Echium amoenum*, *Epicoccum nigrum* from *Rosa canina*, *Fusarium* sp. and *Fusarium equiseti* from *Calendula officinalis* and *Lecanicillium aphanocladii* from *Physalis peruviana*. To screen the phytochemical derivatives of ethyl acetate of these endophytic fungal isolates, the extracts were subjected to phytochemical analysis by gas chromatography-mass spectrometry (GC-MS). GC-MS spectrum of the compounds found in the extracts of the endophytic fungi was matched with the standard compounds present in the WILEY8 library and the National Institute of Standards and Technology (NIST14) library. The GC-MS analysis

of extracts from these endophytic fungi revealed the presence of 49 phytochemicals such as 2,4-Di-tert-butylphenol, Hexadecane, Eicosane, Octadecane, Docosane, Nonadecane, Isopropyl myristate, Hexadecanoic acid, Undecane, Methyl stearate and so on. The results of the present study acknowledge that the endophytic fungi of these medicinal plants are the potential source of biologically active compounds and envisage the possible drug discovery using them. In addition, the compounds 14-Beta-H-Pregna and Cyclohexane, 1, 1'-(2-methyl-1,3-propanediyl) bis are reported here for the first time as fungal metabolites.

Keywords: Endophytes, Fungi, Medicinal plants, Secondary metabolites, GC-MS.

INTRODUCTION

Endophytic fungi are a community of fungi that colonize internal tissues of higher plants, asymptotically (Fouda et al. 2015). Scientific resources consider fungal endophytes as those that live at least part of their life cycle inside diverse plant tissues including leaves, stems, seeds and roots, without any visible damage to their host plants (Fouda et al. 2015). A mutualistic interaction occurs between fungal endophytes and their hosts, the plant provides nutrition and protection for the fungus in one hand, the fungus helps with the competitiveness and growth of the plant by protecting and preserving it against abiotic and biotic stresses on the other hand (Strobel & Daisy 2004). Moreover, plants colonized by endophytes have benefits such as plant growth promotion by helping in the production of growth hormones and secondary metabolites, hence, are generally healthier than those lacking endophytic interaction (Faeth & Fagan 2002). The communication between endophytes and their host plants can also affect the formation of metabolic products in both partners (Faeth & Fagan 2002). It has been revealed that in this mutualistic interactions, fungal endophytes can produce bioactive compounds analogous to plant metabolites and accordingly it has been suggested that, this phenomenon results from co-evolution (Faeth & Fagan 2002). During co-evolution, fungal endophytes gently were

Submitted 4 Dec 2022, accepted for publication 1 March 2023

✉ Corresponding Author: E-mail: m.tajick@sanru.ac.ir

© 2023, Published by the Iranian Mycological Society

<http://mij.areeo.ac.ir>

acclimatized to specific microenvironments using approaches such as the uptake of plant DNA fragments into their genomes or the insertion of their DNA into the host plant genome (Soares et al. 2017). Although many plant species usually host several endophyte species, only a few plants have ever been investigated for their endophytic diversity and biology, therefore, enormous rich potential resources exist for the description of novel fungi (Strobel & Daisy 2004). Some kinds of plant communities such as plants with an ethnobotany background like those used as a medicine, have been selected generally for the isolation of beneficial endophytes (Strobel & Daisy 2004). It has been revealed that endophytic fungi have an affecting role in the quality and quantity of crude drugs through a particular fungus-host interaction in medicinal plants (Faeth & Fagan 2002). Therefore, endophytic fungi are mostly isolated from medicinal herbs to acquire bioactive compounds for remedial activities which are already obtained from host plants (Soares et al. 2017; Musavi et al. 2015).

Plants and fungi are wealthy sources of thousands of bioactive compounds (Musavi et al. 2015). The bioactive compounds resulting from natural resources can be reviewed and evaluated for pharmacological characterization. The triumph of these resources in finding new and effective drug compounds that can be beneficial for humans was considered further with the discovery of biosynthesis of Taxus-derived, anticancer compound Taxol from endophytic fungi *Taxomyces andreanae* Strobel, A. Stierle, D. Stierle & W.M. Hess and *Pestalotiopsis microspora* (Speg.) G.C. Zhao & Nan Li isolated from *Taxus wallichiana* Zucc. (Stierle et al. 1993). Taxol, arguably the most successful anti-cancer drug of all time, is identified and its source, is the bark of the *Taxus* species (Strobel et al. 1996). Capsaicin, an alkaloid of *Capsicum annuum* L. which has anti-inflammatory, thermogenic, anti-lithogenic, gastro-stimulatory, anti-cancer, anti-diabetic and cardio-protective attributes, has also been produced by *Alternaria alternata* (Fr.) Keissl. isolated from *C. annuum* as an endophytic fungus (Musavi et al. 2015). *Huperzia serrata* Thunb., a tropical medicinal moss, is used for the treatment of Alzheimer's disease and its biologically active alkaloid huperzine A (HupA) was also isolated from its endophytic fungus, *Acremonium* sp. (Musavi et al. 2015). A bioactive natural compound Silymarin which has anti-oxidant, anti-hepatitic, cardioprotective, immunomodulatory, anti-inflammatory, anti-metastatic and hepatoprotective activities, found in *Silybum marianum* (L.) Gaertn fruits, has also been reported from the endophytic fungus *Aspergillus iizukae* Sugiy. isolated from this plant (El-Elimat et al. 2014).

In the present research, the seeds of several important medicinal plants including *Calendula officinalis* L., *Echium amoenum* Fisch & Mey, *Physalis peruviana* L. and *Rosa canina* L. were used for the isolation of endophytic fungi. *Calendula*

officinalis, is a medicinal herb that is used in traditional systems of medicine for herpes, scars, ulcers, skin damage and treating wounds, blood purification and frost-bite. Moreover, it is especially used because of its diverse biological activities to treat diseases like gastro-intestinal, gynecological, diabetic, in some cases of burns, analgesic and anti-inflammatory. This plant is wealthy in many pharmaceutical active ingredients like amino acid, flavonoids, glycosides, volatile oil, steroids and carotenoids (Ashwlayan & Verma 2018). *Echium amoenum* is one of the most significant medicinal plants in Iranian traditional medicine and its flowers have been applied as anti-inflammatory, analgesic, demulcent, sedative and anxiolytic in the folk medicine of Iran. The antioxidative stress potential of *E. amoenum* may be due to its bioactive anti-oxidant components, especially rosmarinic acid and flavonoids (Ranjbar et al. 2006). *Physalis peruviana* has a wonderful medicinal value for curing various diseases such as rheumatism, malaria, leukemia, diabetes, asthma, ulcers, hepatitis, dermatitis and cancer (Ramírez et al. 2013). The fruit of *P. peruviana* was exhibited to have both anti-oxidant and anti-inflammatory activities and contains high levels of vitamin A, vitamin B-complex and vitamin C (Pardo et al. 2008). *Rosa canina* contains various vitamins (especially vitamin C), carotenoids, polyphenols, fatty acids and carbohydrates. Moreover, there is evidence of anti-bacterial, anti-diabetic, anti-cancer and anti-obesity properties of this medicinal plant (Selahvarzian et al. 2018). The current study aimed to identify potential bioactive metabolites of five endophytic fungi associated with seeds of the above-mentioned medicinal plants

MATERIALS AND METHODS

Sample collection and isolation of endophytic fungi

Plants and seeds of medicinal plants *C. officinalis*, *E. amoenum*, *P. peruviana* and *R. canina* were collected from Mazandaran Province in Iran during the years 2018–2019 and transferred to the laboratory in sterile Zipper bags in portable cool chambers (4 °C). Endophytic fungal isolations were carried out aseptically from healthy seeds as follows: after primarily washing with tap water, seed samples were processed for surface sterilization with 75% ethanol for 1 min, 3% sodium hypochlorite for 5 min, 70% ethanol for 30 s and, then washed three times with sterile distilled water. Dried plant seeds were cut into 2–3 pieces under sterile conditions and were plated on Petri dishes containing potato dextrose agar (PDA; Merck, Darmstadt, Germany). The plates were incubated at 25 ± 1 °C for 7–12 days and were checked daily. Fungal hyphae and fruiting structures emerging from the plated seeds were subcultured on water agar (WA) media and maintained as the pure culture at 4 °C for further use. The effectiveness of the seed surface sterilization method was evaluated by placing individual seeds onto Petri dishes

containing PDA medium and the plates were incubated at 25 ± 1 °C for 7–12 days.

DNA isolation, PCR amplification and sequencing
Mycelium was scraped from the surface of fungal cultures growing on PDA using a sterile scalpel blade and DNA was extracted according to the protocol of Möller et al. (1992). The DNA samples were diluted 50–100 times in preparation for further DNA amplification reactions. The primers ITS1F (Gardes & Bruns 1993) and ITS4 (White et al. 1990) were used to amplify part of the internal transcribed spacer regions and intervening 5.8S nrRNA gene (ITS) of the nrRNA operon. The polymerase chain reaction (PCR), mixture and amplification condition were performed in a final volume of 25 μ l as explained in Bakhshi & Braun (2022). The PCR products were sequenced with the amplified primers at a commercial sequencing service provided at Microsynth Company (Balgach, Switzerland). Obtained sequences were assembled with MEGA v. X software and consensus sequences were manually generated from the forward and reverse sequences. The consensus regions of sequences were compared with the reference sequences using the BLAST algorithm and submitted to the NCBI GenBank nucleotide database (<http://www.ncbi.nlm.nih.gov/blast>). The obtained sequences from GenBank and the novel generated sequences during this study were aligned and subjected to construct a phylogenetic tree using MrBayes v. 3.2.6 as elucidated by Bakhshi & Braun (2022). *Melampsora caprearum* (DC.) Thüm. (KU550034) was used as the outgroup taxon.

Extraction of metabolites

The metabolites were extracted according to the method of Siddiquee et al. (2012) as follows. The pure fungal colonies were cultured onto 50 ml potato dextrose broth (PDB; Q-LAB) media. After 10 days of incubation, extraction was performed by adding 20 ml Ethyl acetate ($C_4H_8O_2$) to the broth culture in Erlenmeyer flasks and the mixture was incubated at 4 °C for 10 min. Then, flasks were shaken for 30 min at 120 rpm. The Erlenmeyer flasks were placed in a stationary state to form two phases for 60 min. Taken the upper phase was transferred to a clean 15 ml falcon tube. The supernatant was spined for 10 min, with 4000 rpm, at room temperature in the centrifuge. Once again, it was taken in the upper aqueous phase and was transferred to a clean 50 ml falcon tube. Then, the supernatant (metabolites and $C_4H_8O_2$) was separated from the liquid culture and evaporated using a rotary evaporator at 45 °C. The residue was dissolved in one ml methanol (CH_4OH), filtered through a 0.2 μ m syringe filter (Millipore) and stored at 4 °C for 24 h before injection to GC-MS. The chemical composition of the ethyl acetate fractions was analyzed by GC-MS.

GC-MS analysis of the compounds

An Agilent Technologies 7890 A gas chromatograph (GC) connected to a 5975 C inert mass spectrometer (MS) detector (Agilent, USA) was applied for the detection of secondary metabolites from the crude

extract of fungal isolates. Furthermore, an HP-5MS fused silica capillary column (Hewlett-Packard, 30 m* 0.25 mm i.d., 0.25 μ m film, cross-linked to 5% phenyl methyl siloxane stationary phase) was used. The software Chemstation (Hewlett-Packard, version A.01.01) was used to control entire system. Electron impact mass spectra were documented at 70 electron voltage and ultra-high pure He (99.999%) gas was applied as the carrier gas at a flow rate of 1 mL/min. The injection volume was 1 μ l and all the injections were conducted in a split-less mode. The temperature of the injector and detector was 250 and 280 °C, respectively. The column oven temperature was firstly set at 50 °C for 5 min, then enhanced to 260 °C (ramp: 4 °C/min) and held for 5 min (Vandendool & Kratz 1963). The compounds found in the extracts of the fungi were recognized by computer matching their GC-MS spectral patterns with the WILEY8 library and the National Institute of Standards and Technology (NIST14) library.

RESULTS AND DISCUSSION

Identification of endophytic fungal isolates

The isolates were identified with the DNA sequence analysis of the ITS region and based on the ITS phylogenetic tree, were assigned to five species as *Acremonium* sp., *Epicoccum nigrum* Link, *Fusarium equiseti* (Corda) Sacc., *Fusarium* sp. and *Lecanicillium aphanocladii* Zare & W. Gams (Fig. 1). The detailed identification of these endophytic fungi with their Genbank accession numbers are presented in Table 1. There are many studies on endophytic fungi from various plants. This is the first report of endophytic fungi associated with seeds of the medicinal plants *C. officinalis*, *E. amoenum*, *P. peruviana* and *R. canina*.

Identified compounds through GC-MS

According to the results of GC-MS analysis, a total of 49 compounds, including 1-Docosanol, 1-Hexanol 2-ethyl, 2,4-Di-tert-butylphenol, 2-Coumaranone, (2'R)-2-(2'-hydroxypropyl)-4-methoxyl-1,3-benzenediol, 14-Beta-H-Pregna, Adipic acid, Benzenepropanoic acid, Bromotetradecane, Cyclohexane,1,1'-(2-methyl-1,3-propanediyl) bis, Cyclonerolidol, Di-2-ethylhexylphthalate, Di-epi-alpha-cedrene (1H-3a,7-Methanoazulene), Diisooctyl phthalate, Di-n-decylsulfone, Dioxomorpholine, Docosane, Dodecane, Dotriacontane, Eicosane, Heneicosane, Heneicosene, Hentriacontane, Heptacosane, Heptadecane, Heptadecanoic acid, Hexacosane, Hexadecane, Hexadecanoic acid, Isopropyl myristate, Lauric acid, Methyl stearate, N-ethyl-1,3-dithioisindoline, Nonadecane, Octadecane, Octadecene, Pentacosane, Pentadecane, Pentatriacontane, Propanediol, Tetradecane, Tetradecyl ester, Tetracosan, Tetratetracontane, Trichoacorenol, Tridecane, Tritriacontane, Undecane and Zaragozaic acid were identified in this study. Of these 15, 18, 18, 31 and 7 compounds were identified in culture filtrates of the endophytic fungi *Acremonium* sp., *E. nigrum*, *F. equiseti*, *Fusarium* sp.

and *L. aphanocladii* respectively. Some metabolites were produced similarly by several species such as Diisooctyl phthalate and Octadecane and some were produced exclusively in one species like Adipic acid and Dioxomorpholine in *Fusarium* sp., 2-Coumaranone in *F. equiseti*, Cyclonerolidol, 1-Docosanol, Pentatriacontane, Heptadecanoic acid, Trichoacorenol and Octadecene in *Acremonium* sp.

and, 14-Beta-H-Pregna in *E. nigrum*. Details of the various secondary metabolites produced by each fungus are presented in Table S1. The retention time and area of the compounds under the described conditions in the GC-MS section are shown in Tables 2 to 6 and GC-MS chromatograms of the ethyl acetate extract of isolates are presented in Figs. 2 to 6.

Table 1. Identification of endophytic fungi isolated from seeds of some medicinal plants using ITS primers.

Endophytic fungal isolates	Host source	Collection accession	GenBank accession
		number	number
<i>Acremonium</i> sp.	<i>Echium amoenum</i>	AAH5-1	OQ162344
<i>Epicoccum nigrum</i>	<i>Rosa canina</i>	AAH13-3	OQ162360
<i>Fusarium equiseti</i>	<i>Calendula officinalis</i>	AAH9-1	OQ162352
<i>Fusarium</i> sp.	<i>Calendula officinalis</i>	AAH9-2	OQ162353
<i>Lecanicillium aphanocladii</i>	<i>Physalis peruviana</i>	AAH4-1	OQ162343

Table 2. Identification of most abundant bioactive compounds from *Acremonium* sp. extract isolated from *E. amoenum* by GC-MS.

Compounds	R _T	Area	Compounds	R _T	Area
	min	(Maximum)		min	(Maximum)
2,4-Di-tert-butylphenol	28.951	608147	Dotriacontane	39.83 7	806720
Hexadecane	31.374	1943814	Pentatriacontane	40.51 6	558901
Trichoacorenol	33.299	7344440	Eicosane	40.67 3	3121508
Cyclonerolidol	35.302	7344154	Heptadecanoic acid	44.60 0	1812040
Octadecane	35.504	997711	Docosane	46.15 6	1884389
Heneicosane	36.013	446423	Eicosane	50.29 2	2033097
Isopropyl myristate	37.465	6407491	Diisooctyl phthalate	53.33 2	652290683




Fig. 1. Phylogenetic tree inferred by Bayesian analysis of the ITS sequence alignment using MrBayes v.3.2.6. The scale bar indicates 0.05 expected changes per site. The tree was rooted to *Melampsora caprearum* (KU550034). The isolates of this study are shown in bold.

Table 3. Identification of most abundant bioactive compounds from *E. nigrum* extract isolated from *R. canina* by GC-MS.

Compounds	R _T min	Area (Maximum)	Compounds	R _T min	Area (Maximum)
14-. Beta.-H-Pregna	4.128	11135152	Hexadecanoic acid	39.951	3590075
1-Hexanol, 2-ethyl	12.337	84562	Tetradecane	40.516	458115
Hexadecane	31.379	1061704	10-Heneicosene	41.943	595219
Tridecane	34.404	548231	Methyl stearate	44.610	1672585
Octadecane	36.775	2824435	Docosane	44.776	1202468
Dodecane	36.947	1125497	Pentacosane	46.167	1288079
Isopropyl myristate	37.471	3916314	Heptadecane	49.280	652177
Eicosane	39.847	1204447	N-ethyl-1,3-dithioisindoline	53.410	886316

Table 4. Identification of most abundant bioactive compounds from *F. equiseti* extract isolated from *C. officinalis* by GC-MS.



Compounds	R _T min	Area (Maximum)	Compounds	R _T min	Area (Maximum)
2-Coumaranone	19.974	981915	Hexadecanoic acid	39.946	3862236
Hexadecane	31.374	2428833	Undecane	40.516	615460
Docosane	34.399	561295	Tetracosane	40.854	898019
Octadecane	35.514	850768	Heptacosane	40.999	520550
Heptadecane, 3-methyl-	36.018	611280	Eicosane	41.679	3437782
Nonadecane	36.775	4951791	Methyl stearate	44.600	1827009
Cyclohexane,1,1'-(2-methyl-1,3-propanediyl) bis	36.936	1227276	Heneicosane	44.771	1264392
Isopropyl myristate	37.471	6735356	Hexacosane	46.161	2011737
Hentriacontane	39.842	1022517	Diisooctyl phthalate	50.021	351547623

Table 5. Identification of most abundant bioactive compounds from *Fusarium* sp. extract isolated from *C. officinalis* by GC-MS.

Compounds	R _T min	Area (Maximum)	Compounds	R _T min	Area (Maximum)
2,4-Di-tert-butylphenol	28.951	1417127	Bromotetradecane	39.847	849959
Dioxomorpholine	29.957	14833923	Hexadecanoic acid	39.942	6833887
Dioxomorpholine	30.990	1697320	Benzenepropanoic acid	40.340	1182369
Hexadecane	31.369	3458348	Heneicosane	40.345	602905
Adipic acid	32.106	3896169	Dotriacontane	40.859	1323393
Di-epi-alpha-cedrene	32.246	1580966	Heptacosane	40.994	219254
Zaragozic acid	32.775	637333	Eicosane	41.673	3632902
Di-n-decylsulfone	34.155	662795	Tetratetracontane	41.004	510019
Docosane	34.394	1172734	Tetradecyl ester	41.674	3632902
Hentriacontane	35.317	1073510	Octadecene	41.943	929945
Pentadecane	35.509	1345791	Isobutyl methyl phthalate	42.467	2904871
Nonadecane	36.771	5485589	Hentriacontane	44.776	2317927
n-Tetracosanol-1	36.936	1602275	Heptadecane	46.162	2127231
Propanediol	36.941	1343047	Heneicosane	49.275	872730
Isopropyl myristate	37.465	7793088	Tritriacontane	50.297	1942157
Lauric acid	38.373	5957526	Diisooctyl phthalate	53.283	484429839

Table 6. Identification of most abundant bioactive compounds from *L. aphanocladii* extract isolated from *P. peruviana* by GC–MS.

Compounds	R _T min	Area (Maximum)	Compounds	R _T min	Area (Maximum)
Hexadecane	31.379	3077182	Methyl stearate	44.605	3332974
Octadecane	36.775	4673430	Docosane	46.167	1867239
Isopropyl myristate	37.476	6204609	Hexatriacontane	50.297	1854631
Hexadecanoic acid, methyl ester	39.946	6263271	Diisooctyl phthalate	53.275	414283706

**Fig. 2.** Chromatogram of the active extract from *Acremonium* sp. showing major volatile compounds by GC–MS.**Fig. 3.** Chromatogram of the active extract from *E. nigrum* showing major volatile compounds by GC–MS.




Fig. 4. Chromatogram of the active extract from *F. equiseti* showing major volatile compounds by GC-MS.




Fig. 5. Chromatogram of the active extract from *Fusarium* sp. showing major volatile compounds by GC-MS.




Fig. 6. Chromatogram of the active extract from *L. aphanocladii* showing major volatile compounds by GC-MS

Potential biological activities of compounds identified in this study

A diverse range of biological activities has been reported in the literature for different compounds identified in this research. To give some examples, Hexadecanoic acid which was produced by all the fungal endophytes studied in this research has been confirmed that displays a wide range of biological activities such as antifungal, antioxidants, hypocholesterolemic prevention and anticancer effects (Anisha & Radhakrishnan 2017). This metabolite was mainly found in the endophytic fungus *Fusarium* sp. obtained from the well-known medicinal plant *Zingiber officinale* Roscoe which is commonly used in traditional medicine (Anisha & Radhakrishnan 2017). Diisooctyl phthalate is another compound that was produced by all the fungal endophytes studied here. Abdel-Motaal et al. (2022), stated that analysis of GC-MS of extracts from three fungal endophytes including *Aspergillus terreus* Thom., *Fusarium solani* (Mart.) Sacc. and *Penicillium verrucosum* Dierckx, isolated from the medicinal plant *Solenostemma argel* (Delile) Hayne., indicated the presence of Diisooctyl phthalate as the main compound. Kalavathi et al. (2017) documented that Diisooctyl phthalate derived from an endophytic fungus, *Penicillium* sp., isolated from the medicinal plant *Tabebuia argentea* (Bureau & K.Schum.) Britton, displayed anti-diabetic activity. Isopropyl myristate which was produced in this study exclusively by *Fusarium* sp. isolated from *Calendula officinalis* has been shown that depicts cytotoxic activity against different cancer cell lines (Mo & Lim 2005). Pelo et al. (2021) declared that *Aureobasidium pullulans* (de Bary & Löwenthal) G. Arnaud, *Cladosporium* sp., *Fusarium* sp., *Hyalodendriella* sp., *Penicillium chrysogenum* Thom. and, *Phialophora* sp. which isolated from the medicinal plant *Solanum mauritianum* Scop. as endophytic fungi, can produce Isopropyl myristate. To illustrate more examples, some of the biological activities reported for all the compounds identified in this study along with some examples of other fungi producing similarly these compounds are listed in Table S1.

To the best of our knowledge, two compounds including 14-Beta-H-Pregna produced exclusively by *E. nigrum* and Cyclohexane,1,1'-(2-methyl-1,3-propanediyl) bis produced exclusively by *F. equiseti* were reported in this research for the first time as fungal metabolites. Previous phytochemical studies on volatile oils of medicinal plants have revealed the production of 14-Beta-H-Pregna by plants (Gharari et al. 2019). In addition, several reports confirmed the biological activities of 14-Beta-H-Pregna which exhibits antibacterial, antifungal and antioxidant effects (Gharari et al. 2019). Furthermore, previous reports confirmed the production of the compound Cyclohexane,1,1'-(2-methyl-1,3-propanediyl) bis by plants and its biological activities as an antioxidant (Delaram et al. 2019). Therefore, these data more

confirm that endophytic fungi are a rich source of novel and bioactive secondary metabolites.

Conclusion

In conclusion, our results reveal the potential of the medicinal plants *C. officinalis*, *E. amoenum*, *P. peruviana* and *R. canina* as a host of important fungal endophytes which possess a precious source of bioactive compounds which have diverse biological activities, such as antioxidant, insecticidal, antimicrobial and anticancer activities. These compounds can increase the tolerance of host plants to biotic and abiotic stresses. Moreover, they can be used for the treatment of diseases and for manufacturing new drugs. Furthermore, these compounds can also be applied in dyeing, polyester, resin, biofuels and cosmetics industries. Therefore, further investigation and research will be required on the structural features and application strategies of these potent resources and also the influence of environmental factors when used in different conditions.

REFERENCES

- Abdel-Motaal FF, Kamel NM, El-Sayed MA, Mohamed Abou-Ellail M. 2022. Biocontrol of okra-rot-causing *Cochliobolus spicifer*-CSN-20 using secondary metabolites of endophytic fungi associated with *Solenostemma argel*. *Annals of Agricultural Sciences* 67(1): 24–33.
- Adongbete EM. 2022. Bioprocessing of pharmacologically important compounds from edible mushrooms in Nigeria. *Arboricultural Association, World Fungi Day, International Seminar*. <https://www.trees.org.uk/Training-Events/World-Fungi-Day>.
- Ahamed A, Ahring BK. 2011. Production of hydrocarbon compounds by endophytic fungi *Gliocladium* species grown on cellulose. *Bioresource Technology* 102(20): 9718–9722.
- Albratty M, Alhazmi HA, Meraya AM, Najmi A, Alam MS, Rehman Z, Moni SS. 2021. Spectral analysis and Antibacterial activity of the bioactive principles of *Sargassum tenerrimum* J. Agardh collected from the Red sea, Jazan, Kingdom of Saudi Arabia. *Brazilian Journal of Biology* 83: 1–10.
- Anisha C, Radhakrishnan EK. 2017. Metabolite analysis of endophytic fungi from cultivars of *Zingiber officinale* Rosc. identifies myriad of bioactive compounds including tyrosol. *3 Biotech* 7(2): 1–10.
- Ashwlayan VDKA, Verma M. 2018. Therapeutic potential of *Calendula officinalis*. *Pharmacy and Pharmacology International Journal* 6(2): 149–155.
- Asong JA, Amoo SO, McGaw LJ, Nkadimeng SM, Aremu AO, Otang-Mbeng W. 2019. Antimicrobial activity, antioxidant potential, cytotoxicity and phytochemical profiling of four plants locally used against skin diseases. *Plants* 8(9): 1–19.

- Bakhshi M, Braun U. 2022. *Acericercospora hyrcanica* gen. et sp. nov. (Mycosphaerellaceae) and *Paramycocentrospora acericola* gen. et sp. nov. (Dothidotthiaceae) on maple trees in Hyrcanian forests. *Mycological Progress* 21(8): 1–15.
- Bhatnagar VS, Bandyopadhyay P, Rajacharya GH, Sarkar S, Poluri KM, Kumar S. 2019. Amelioration of biomass and lipid in marine alga by an endophytic fungus *Piriformospora indica*. *Biotechnology for Biofuels* 12(1): 1–9.
- Bhimba BV, Franco DA, Jose GM, Mathew JM, Joel EL. 2011. Characterization of cytotoxic compound from mangrove derived fungi *Irpex hydroides* VB4. *Asian Pacific Journal of Tropical Biomedicine* 1(3): 223–226.
- Chowdhary K, Kaushik N. 2018. Biodiversity study and potential of fungal endophytes of peppermint and effect of their extract on chickpea rot pathogens. *Archives of Phytopathology and Plant Protection* 51: 139–155.
- Das M, Prakash HS, Nalini MS. 2020. Antibacterial metabolites from *Bipolaris specifera*, an endophytic fungus from the endemic medicinal plant, *Zingiber nimmonii* (J. Graham) Dalzell. *3 Biotech* 10(7): 1–8.
- Delaram J, Esmaeilzadeh Bahabadi S, Ijbari HA. 2019. Study on phytochemical composition and antioxidant activity of different organs of 3 species of *Chenopodium* in Sistan region. *Journal of Plant Research (Iranian Journal of Biology)* 32(3): 570–582.
- Dhankhar S, Yadav JP. 2013. Investigating antimicrobial properties of endophytic fungi associated with *Salvadora oleoides* decne. *Anti-Infective Agents in Medicinal Chemistry* 11(1): 48–58.
- El-Amrani M, Debbab A, Aly AH, Wray V, Dobretsov S, Müller WEG, Lin W, Lai D, Proksch P. 2012. Farinomalain derivatives from an unidentified endophytic fungus isolated from the mangrove plant *Avicennia marina*. *Tetrahedron Letters* 53(49): 6721–6724.
- El-Elimat T, Raja HA, Graf TN, Faeth SH, Cech NB, Oberlies NH. 2014. Flavonolignans from *Aspergillus iizukae*, a fungal endophyte of milk thistle (*Silybum marianum*). *Journal of Natural Products* 77: 193–199.
- El-Zawawy N, Metwally M, El-Salam A. 2020. Antitumor and antimicrobial activities of endophytic fungi obtained from Egyptian *Urospermum picroides*. *International Journal of Cancer and Biomedical Research* 4(3): 187–199.
- Faeth SH, Fagan WF. 2002. Fungal endophytes: common host plant symbionts but uncommon mutualists. *Integrative and Comparative Biology* 42: 360–368.
- Fouda AH, Hassan SED, Eid AM, Ewais EED. 2015. Biotechnological applications of fungal endophytes associated with medicinal plant *Asclepias sinaica* (Bioss.). *Annals of Agricultural Science* 60(1): 95–104.
- Gardes M, Bruns TD. 1993. ITS primers with enhanced specificity for basidiomycetes-application to the identification of mycorrhizae and rusts. *Molecular Ecology* 2(2): 113–118.
- Gharari Z, Bagheri Kh, Danafar H, Sharafi A. 2019. Essential oil composition of two *Scutellaria* species from Iran. *Journal of Traditional Chinese Medical Sciences* 6: 244–253.
- Hegazy MMS, El-Hela AA, Ismail SK. 2015. Isolation and structure elucidation of bioactive secondary metabolites from *Kalanchoe tomentosa* and some of its endophytic fungi. Thesis of the requirements of the Ph.D. Degree In Pharmaceutical Sciences, Pharmacognosy Department Faculty of Pharmacy of Al-Azhar University, Nasr City, Cairo: p.327.
- Huang L, Zhu X, Zhou SH, Cheng Z, Shi K, Zhang C, Shao H. 2021. Phthalic acid esters: natural sources and biological activities. *Toxins* 13(7): 1–17.
- Jayaram H, Marigowda V, Saraswathi KJT. 2021. Secondary metabolite production and terpenoid biosynthesis in endophytic fungi *Cladosporium cladosporioides* isolated from wild *Cymbopogon martinii* (Roxb.) Wats. *Microbiological Research* 12: 812–828.
- Kalavathi MK, Chandrappa CP, Channabasava R, Ramachandra YL, Padmalatha RS, Ravishankar RV, Govindappa M. 2017. Anti-diabetic activity of endophytic fungi, *Penicillium* species of *Tabebuia argentea*; in silico and experimental analysis. *Research Journal of Phytochemistry* 11(2): 90–110.
- Kandasamy S, Sahu SK, Kandasamy K. 2012. In silico studies on fungal metabolite against skin cancer protein (4,5-diarylisoazole hsp90 chaperone). *International Scholarly Research Network, Dermatology* 3. <https://doi.org/10.5402/2012/626214>
- Kanjana M, Kanimozhi G, Udayakumar R, Panneerselvam A. 2019. GC-MS analysis of bioactive compounds of endophytic fungi *Chaetomium globosum*, *Cladosporium tenuissimum* and *Penicillium janthinellum*. *Journal of Biomedical and Pharmaceutical Sciences* 2(2): 1–10.
- Kaur N, Arora DS, Kalia N, Kaur M. 2020. Bioactive potential of endophytic fungus *Chaetomium globosum* and GC-MS analysis of its responsible components. *Scientific Reports* 10(1): 1–10.
- Khiralla A, Spina R, Varbanov M, Philippot S, Lemièrre P, Slezacek-Deschaumes S, André P, Mohamed I, Mohamed Yagi S, Laurain-Mattar D. 2020. Evaluation of antiviral, antibacterial and antiproliferative activities of the endophytic fungus *Curvularia papendorffii*, and isolation of a new polyhydroxyacid. *Microorganisms* 8(9): 1–21.
- Kouipou Toghueo RM, Eugénie Kemgne AM, Eke P, Mbekou Kanko MI, Dize D, Sahal D, Boyom FF.

2019. Antiplasmodial potential and GC-MS fingerprint of endophytic fungal extracts derived from Cameroonian *Annona muricata*. *Journal of Ethnopharmacology* 235: 111–121.
- Kumari N, Menghani E, Mithal R. 2019. Bioactive compounds characterization and antibacterial potentials of actinomycetes isolated from rhizospheric soil. *Journal of Scientific and Industrial Research* 78: 793–798.
- Li Y, Kumar PS, Tanb Q, Tana X, Yuana M, Luoc J, Hea M. 2021. Diversity and chemical fingerprinting of endo-metabolomes from endophytes associated with *Ampelopsis grossedentata* (Hand.-Mazz.) W. T. Wang possessing antibacterial activity against multidrug resistant bacterial pathogens. *Journal of Infection and Public Health* 14: 1917–1926.
- Luo YP, Zheng CJ, Chen GY, Song XP, Wang Z. 2019. Three new polyketides from a mangrove-derived fungus *Colletotrichum gloeosporioides*. *The Journal of Antibiotics* 72: 513–517.
- Marrufu T, Nazzaro F, Mancini E, Fratianni F, Coppola R, De Martino L, Agostinho AB, De Feo V. 2013. Chemical composition and biological activity of the essential oil from leaves of *Moringa oleifera* Lam. cultivated in Mozambique. *Molecules* 18(9): 10989–11000.
- Meshram V, Kapoor N, Chopra G, Saxena S. 2017. *Muscodor camphora*, a new endophytic species from *Cinnamomum camphora*. *Mycosphere* 8(4): 568–582.
- Mishra VK, Passari AK, Chandra P, Leo VV, Kumar B, Uthandi S, Thankappan S, Gupta VK, Singh BP. 2017. Determination and production of antimicrobial compounds by *Aspergillus clavatonanicus* strain MJ31, an endophytic fungus from *Mirabilis jalapa* L. using UPLC-ESI-MS/MS and TD-GC-MS analysis. *Plos One* 12(10): 1–24.
- Mo Y, Lim LY. 2005. Preparation and in vitro anticancer activity of wheat germ agglutinin (WGA)-conjugated PLGA nanoparticles loaded with paclitaxel and isopropyl myristate. *Journal of Controlled Release* 107: 30–42.
- Möller EM, Bahnweg G, Sandermann H, Geiger HH. 1992. Simple and efficient protocol for isolation of high molecular weight DNA from filamentous fungi, fruit bodies, and infected plant tissues. *Nucleic Acids Research* 20(22): 6115–6116.
- Musavi SF, Dhavale A, Balakrishnan RM. 2015. Optimization and kinetic modeling of cell-associated camptothecin production from an endophytic *Fusarium oxysporum* NFX06. *Preparative Biochemistry and Biotechnology* 45: 158–172.
- Pardo JM, Fontanilla MR, Ospina LF, Espinosa L. 2008. Determining the pharmacological activity of *Physalis peruviana* fruit juice on rabbit eyes and fibroblast primary cultures. *Investigative Ophthalmology and Visual Science* 49: 3074–3079.
- Pelo SP, Adebo OA, Green E. 2021. Chemotaxonomic profiling of fungal endophytes of *Solanum mauritanium* (alien weed) using gas chromatography high resolution time-of-flight mass spectrometry (GC-HRTOF-MS). *Metabolomics* 17(5): 1–13.
- Pinto ME, Araujo SG, Morais MI, Sa NP, Lima CM, Rosa CA, Siqueira EP, Johann S, Lima LA. 2017. Antifungal and antioxidant activity of fatty acid methyl esters from vegetable oils. *Anais da Academia Brasileira de Ciências* 89: 1671–1681.
- Raimundo K, de Campos Bortolucci W, Luiz Rahal I, Marko de Oliveira HL, Piau Junior R, de Araujo Almeida Campo CF, Eduardo Gonçalves J, Andrea Linde G, de Barros Colauto N, Cristiani Gazim Z. 2021. Insecticidal activity of *Gallesia integrifolia* (Phytolaccaceae) essential oil. *Boletín Latinoamericano y del Caribe de Plantas Medicinales y Aromáticas* 20(1): 38–50.
- Ramírez F, Fischer G, Davenport TL, Pinzón JCA, Ulrichs C. 2013. Cape gooseberry (*Physalis peruviana* L.) phenology according to the BBCH phenological scale. *Scientia Horticulturae* 162: 39–42.
- Ranjbar A, Khorami S, Safarabadi M, Shahmoradi A, Malekirad AA, Vakilian K, Mandegary A, Abdollahi M. 2006. Antioxidant activity of Iranian *Echium amoenum* Fisch & CA Mey flower decoction in humans: a cross-sectional before/after clinical trial. *Evidence-Based Complementary and Alternative Medicine* 3(4): 469–473.
- Robey MT, Ye R, Bok JW, Clevenger KD, Islam MN, Chen C, Gupta R, Swyers M, Wu E, Gao P, Thomas PM, Wu CC, Keller NP, Kelleher NL. 2018. Identification of the first diketomorpholine biosynthetic pathway using FAC-MS technology. *ACS Chemical Biology* 13(5): 1142–1147.
- Sarsaiya S, Jain A, Fan X, Jia Q, Xu Q, Shu F, Zhou Q, Shi J, Chen J. 2020. New insights into detection of a dendrobine compound from a novel endophytic *Trichoderma longibrachiatum* strain and its toxicity against phytopathogenic bacteria. *Frontiers in Microbiology* 11: 1–12.
- Sawant A, Rodrigues BF, Sardesai Y. 2018. Antimicrobial and anti-cancer activity of *Setosphaeria monoceras*, an endophytic fungus associated with *Tropical mangrove* plant. *World Journal of Pharmacy and Pharmaceutical Sciences* 7(6): 1315–1329.
- Saxena S, Strobel GA. 2021. Marvellous *Muscodor* spp.: update on their biology and applications. *Microbial Ecology* 82: 5–20.
- Selahvarzian A, Alizadeh A, Baharvand PA, Eldahshan OA, Rasoulzadeh B. 2018. Medicinal properties of *Rosa canina* L. *Herbal Medicines Journal* 3(1): 77–84.
- Siddiquee S, Cheong BE, Khanam T, Hossain K. 2012. Separation and identification of volatile compounds from liquid cultures of *Trichoderma harzianum* by GC-MS using three different

- capillary columns. *Journal of Chromatographic Science* 50(4): 358–368.
- Soares DA, Rosa LH, Silva JFM, Pimenta RS. 2017. A review of bioactive compounds produced by endophytic fungi associated with medicinal plants. *Boletim do Museu Paraense Emílio Goeldi Ciências Naturais* 12(3): 331–352.
- Sridharan AP, Thangappan S, Karthikeyan G, Nakkeeran S, Uthandi S. 2021. Metabolites of *Trichoderma longibrachiatum* EF5 inhibits soil borne pathogen, *Macrophomina phaseolina* by triggering amino sugar metabolism. *Microbial Pathogenesis* 150: 1–9.
- Stierle A, Strobel G, Stierle D. 1993. Taxol and Taxane production by *Taxomyces andreanae*, an endophytic fungus of Pacific Yew. *Science* 260: 214–216.
- Strobel GA, Hess WM, Ford E, Sidhu RS, Yang X. 1996. Taxol from fungal endophytes and the issue of biodiversity. *Journal of Industrial Microbiology and Biotechnology* 17: 417–423.
- Strobel G, Daisy B, Castillo U, Harper J. 2004. Natural Products from Endophytic Microorganisms. *Journal of Natural Products* 67(2): 257–268.
- Sun Y, Wang C, Du G, Deng W, Yang H, Li R, Xu Q, Guo Q. 2022. Two nematicidal compounds from *Lysinimonas* M4 against the pine wood nematode, *Bursaphelenchus xylophilus*. *Forests* 13(8): 1–14.
- Tanapichatsakul C, Khruengsai S, Monggoot S, Pripdeevech P. 2019. Production of eugenol from fungal endophytes *Neopestalotiopsis* sp. and *Diaporthe* sp. isolated from *Cinnamomum loureiroi* leaves. *PeerJ* 7: 1–21.
- Thelma J, Balasubramanian C. 2021. Screening and bioactivity profiling of marine *Bacillus* species against human pathogens. *Journal of Scientific Research* 65(5): 72–95.
- Vandendool H, Kratz PD. 1963. A generalization of the retention index system including linear temperature programmed gas-liquid partition chromatography. *Journal of Chromatography A* 11: 463–471.
- White TJ, Bruns TD, Lee SB, Taylor JW. 1990. Amplification and direct sequencing of fungal ribosomal RNA Genes for phylogenetics. In Book: *A Guide to Methods and Applications*, New York :315–322.
- Wu SH, Zhao LX, Chena YW, Huang R, Miaoa CP, Wang J. 2011. Sesquiterpenoids from the endophytic fungus *Trichoderma* sp. pr-35 of *Paeonia delavayi*. *Chemistry and Biodiversity* 8: 1717–1723.
- Zhang M, Zhao J, Liu JM, Chen R, Xie KB, Chen DW, Feng KP, Zhang D, Dai J. 2017. Neural anti-inflammatory sesquiterpenoids from the endophytic fungus *Trichoderma* sp. Xy24. *Journal of Asian Natural Products Research* 19(7): 651–658.
- Zhang Y, Zhao J, Wang J, Shan T, Mou Y, Zhou L, Wang J. 2011. Chemical composition and antimicrobial activity of the volatile Oil from *Fusarium tricinctum*, the endophytic fungus in *Paris polyphylla* var. *yunnanensis*. *Natural Product Communications* 6(11): 1759–1762.
- Zhao F, Wang P, Su Z, Li S. 2020. Natural sources and bioactivities of 2,4-di-tert-butylphenol and its analogs. *National Center for Pharmaceutical Crops. Toxins* 12(1): 1–26.
- Zhao J, Shan T, Huang Y, Liu X, Gao X, Wang M, Jiang W, Zhou L. 2009. Chemical composition and in vitro antimicrobial activity of the volatile oils from *Gliomastix murorum* and *Pichia guilliermondii*, two endophytic fungi in *Paris polyphylla* var. *yunnanensis*. *Natural Product Communications* 4(11): 1491–1496.

بررسی ترکیبات زیست فعال عصاره قارچ‌های اندوفیت جدا شده از بذور چند گیاه دارویی با روش کروماتوگرافی گازی و طیف سنجی جرمی (GC-MS)

عارفه اصغری^۱، محمدعلی تاجیک قنبری^۱، مونس بخشی^۲ ولی‌الله بابایی‌زاد^۱

۱- گروه گیاهپزشکی، دانشکده علوم زراعی، دانشگاه علوم کشاورزی و منابع طبیعی ساری، ساری، ایران.

۲- بخش تحقیقات رستنی‌ها، مؤسسه تحقیقات گیاهپزشکی کشور، سازمان تحقیقات، آموزش و ترویج کشاورزی، تهران، ایران.

چکیده: قارچ‌های اندوفیت گروهی از جوامع قارچی مرتبط با میزبان هستند و در عین حال برای میزبان خود سودمند می‌باشند. قارچ‌های اندوفیت گیاهی با توجه به شرایط محیط زندگی ویژه خود، گروه‌های متنوعی از متابولیت‌های زیست فعال با ویژگی‌های ساختاری متفاوت را تولید می‌کنند. ترکیبات زیست فعال تولید شده توسط قارچ‌های اندوفیت، از اثرات قابل توجهی برای افزایش سازگاری آنها و گیاهان میزبان-شان، از قبیل تحمل به تنش‌های زنده و غیرزنده برخوردارند. علاوه بر این، برخی از این متابولیت‌ها دارای اهمیت دارویی و زیست محیطی نیز هستند. در تحقیق حاضر پنج قارچ اندوفیت از بذر تعدادی از گیاهان دارویی جداسازی شد. جدایه‌های به دست آمده بر اساس داده‌های توالی نوکلئوتیدی ناحیه ژنومی ITS تحت آرایه‌های *Acremonium* sp. از بذر گل گاوزبان (*Echium amoenum*)، *Epicoccum nigrum* و از بذر نسترن کوهی (*Rosa canina*)، *Fusarium* sp. و *Fusarium equiseti* از بذر همیشه بهار (*Calendula officinalis*) و از بذر نسترن کوهی *Lecanicillium aphanocladii* از بذر عروسک پشت پرده (*Physalis peruviana*) شناسایی شدند. به منظور غربالگری ترکیبات فیتوشیمیایی عصاره اتیل استات این جدایه‌های قارچی اندوفیت، عصاره‌ها با استفاده از کروماتوگرافی گازی-طیف سنجی جرمی (GC-MS)، مورد آزمایش فیتوشیمیایی قرار گرفتند. طیف GC-MS عصاره اتیل استات جدایه‌های قارچی، با ترکیبات استاندارد موجود در WILEY8 و NIST14 مطابقت داده شد. تجزیه و تحلیل این داده‌ها، وجود ۴۹ ترکیب گیاهی از قبیل Hexadecane, 2,4-Di-tert-butylphenol, Methyl Undecane, Hexadecanoic acid, Isopropyl myristate, Nonadecane, Docosane, Octadecane, Eicosane و ... را در این جدایه‌ها آشکار نمود. نتایج این مطالعه تایید کرد که قارچ‌های اندوفیت جدا شده از گیاهان دارویی، منبع بالقوه‌ای از ترکیبات دارای خواص بیولوژیکی هستند. علاوه بر این، دو ترکیب 14-Beta-H-Pregna و Cyclohexane,1,1'-(2-methyl-1,3- propanediyl) bis برای اولین بار در این مطالعه به عنوان متابولیت‌های قارچی معرفی می‌شوند.

کلمات کلیدی: اندوفیت‌ها، قارچ‌ها، گیاهان دارویی، متابولیت‌های ثانویه، GC-MS.